

DEPARTMENT OF AGRICULTURE AND FISHERIES, SOUTH AUSTRALIA

Agronomy Branch Report

BOTANICAL COMPOSITION AND PRODUCTIVITY OF FIVE
GRAZED PASTURES ON KANGAROO ISLAND

P.R. Gibson

Senior Research Officer (Agronomy)

Report No. 88

BOTANICAL COMPOSITION AND PRODUCTIVITY
OF FIVE GRAZED PASTURES ON KANGAROO
ISLAND

1977

P.R. GIBSON

SENIOR RESEARCH OFFICER (AGRONOMY)

TABLE OF CONTENTS

	<u>Page</u>
List of Figures	v
List of Tables	vi
List of Appendix Tables	vii
Summary	ix
1. INTRODUCTION	1
2. ACKNOWLEDGEMENTS	3
3. BACKGROUND TO EXPERIMENTAL PROGRAM	4
4. AIM OF EXPERIMENT	6
5. ENVIRONMENT	7
6. SITE OF EXPERIMENT	11
7. EXPERIMENTAL METHODS	13
7.1 Site preparation	13
7.2 Treatments	13
7.2.1 Pasture treatments	13
7.2.2 Stocking rates	15
7.2.3 Layout and allocation of treatments	15
7.3 Pasture establishment	17
7.4 Fertilizer application	17
7.5 Selection and allocation of experimental animals	18
7.6 Grazing management of pastures in 1970	19
7.7 Routine management	19
7.7.1 Pastures	19
7.7.2 Animals	20
7.8 Data collection procedures	21
7.8.1 Pasture data 1970	21
7.8.2 Pasture data 1971-1974	21
7.8.3 Subterranean clover seed	23

TABLE OF CONTENTS (Contd.)

	<u>Page</u>	
7.8.4	Sheep live weights	24
7.8.5	Annual wool production	24
7.8.6	Fleece characteristics	25
7.8.7	Seasonal wool production	25
7.9	Data analysis	26
8.	RESULTS	28
8.1	Pasture establishment	28
8.2	Pasture production 1970	28
8.3	Botanical composition 1971-1974	29
8.3.1	Sown grasses	29
8.3.2	Subterranean clover	35
8.3.3	Volunteer annual species	37
8.4	Growth, crude utilization and availability of pasture	46
8.4.1	1971 Season	46
8.4.2	1972 Season	50
8.4.3	1973 and 1974 Seasons	51
8.5	Subterranean clover seed production and reserves	52
8.6	Sheep liveweight data	57
8.6.1	General trends	57
8.6.2	1971	58
8.6.3	1972 and 1973	59
8.6.4	1974	60
8.7	Annual wool production	62
8.8	Fleece characteristics	66
8.9	Seasonal wool production	66
9.	DISCUSSION AND CONCLUSION	72
9.1	Animal production in relation to botanical composition	72
9.1.1	The importance of subterranean clover	72
9.1.2	The role of other species	75

<u>TABLE OF CONTENTS (Contd.)</u>		<u>Page</u>
9.2	Persistence of sown species	76
9.2.1	Grasses	76
9.2.2	Subterranean clover	78
9.3	Invasion of non-leguminous species	81
9.4	Interaction of stocking rate and botanical composition in the determination of animal production	82
9.5	Wool production and fleece characteristics	83
9.6	Efficiency of high stocking rates	85
9.7	Conclusion	86
10.	APPENDICES	88
11.	BIBLIOGRAPHY	117

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
-	Plan of experimental paddocks	16
1	Changes in botanical composition with time and stocking rate	32
2	Growth, crude utilization and availability of pasture	47
3	Changes in live weights and regression coefficients with time	61
4	Relation of wool produced per head to stocking rate	64
5	Relation of wool produced per hectare to stocking rate	65
6	Changes in wool growth rates and regression coefficients with time	71

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	List of scientific and common names of pasture species	9
2	Rainfall recorded at the Kangaroo Island Research Centre	10
3	Purity, viability and establishment of sown pasture seed	30
4	Growth and availability of pasture in 1970	30
5	Proportion of sown grass in the pasture (1971-1974)	33
6	Density of perennial plants in June of each year	34
7	Density of clover plants in June of each year	39
8	Proportion of Yarloop in the total clover component (1971 and 1974)	39
9	Proportion of clover in the pasture (1971-1972)	40
10	Proportion of clover in the pasture (1973-1974)	41
11	Proportion of broad-leaved volunteers in the pasture (1971-1972)	42
12	Proportion of broad-leaved volunteers in the pasture (1973-1974)	43
13	Proportion of narrow-leaved volunteers in the pasture (1971-1972)	44
14	Proportion of narrow-leaved volunteers in the pasture (1973-1974)	45
15	Annual pasture production for the experimental period	48
16	Clean seed yields of subterranean clover	53
17	Availability of subterranean clover (1971-1972)	54
18	Availability of subterranean clover (1973-1974)	55
19	Annual clean wool production	63
20	Fleece characteristics (1971-1972)	69
21	Fleece characteristics (1973-1974)	70
22	Efficiency of wool production in relation to pasture production and rainfall	87

LIST OF APPENDIX TABLES

<u>Appendix Table</u>	<u>Page</u>
1 Meteorological data from the K.I.R.C. (1970)	88
2 Meteorological data from the K.I.R.C. (1971)	89
3 Meteorological data from the K.I.R.C. (1972)	90
4 Meteorological data from the K.I.R.C. (1973)	91
5 Meteorological data from the K.I.R.C. (1974)	92
6 Pasture production in 1971	93
7 Pasture production in 1972	94
8 Pasture production in 1973	95
9 Pasture production in 1974	96
10 Crude utilization of pasture in 1971	97
11 Crude utilization of pasture in 1972	98
12 Crude utilization of pasture in 1973	99
13 Crude utilization of pasture in 1974	100
14 Availability of pasture in 1971-1972	101
15 Availability of pasture in 1973-1974	102
16 Mean live weights of wethers in 1970-1971	103
17 Mean live weights of wethers in 1971-1972	104
18 Mean live weights of wethers in 1972-1973	105
19 Mean live weights of wethers in 1973-1974	106
20 Mean live weights of wethers in 1974-1975	107
21 Regression coefficients (L.W. on S.R.) in 1971-1973	108
22 Regression coefficients (L.W. on S.R.) in 1973-1975	109
23 Growth rate of clean wool in 1971-1972	110
24 Growth rate of clean wool in 1972-1973	111
25 Growth rate of clean wool in 1973-1974	112
26 Growth rate of clean wool in 1974-1975	113

LIST OF APPENDIX TABLES (Contd.)

<u>Appendix Table</u>		<u>Page</u>
27	Regression coefficients (Wool growth rate on S.R.) in 1971-1973	114
28	Regression coefficients (Wool growth rate on S.R.) in 1973-1975	115
29	Number and cause of deaths of experimental wethers	116

SUMMARY

A stocking rate experiment evaluating five pasture types was undertaken in a Mediterranean-type climate on the Kangaroo Island Research Centre, Parndana East, South Australia.

The hypothesis tested was that the renovation of an existing volunteer annual grass - Yarloop subterranean clover (*Trifolium subterraneum* L. cv. Yarloop) pasture by tillage, seedbed preparation and sowing of a mixture of the best available grass species and low oestrogenic subterranean clovers, would result in increased live weight and wool production from grazing wethers.

Merino wethers were continuously grazed for four years at six rates of stocking (10, 11, 13, 14, 15 and 17 sheep ha⁻¹) on each of the following five pasture treatments.

Treatment (A) - An unrenovated, volunteer annual grass-subterranean clover pasture dominated by Yarloop subterranean clover and barley grass (*Hordeum leporinum* Link.).

Treatments B, C, D and E - Tillage, seed bed preparation and sowing of grass, together with Mt. Barker and Woogenellup subterranean clovers. Treatment grasses sown were as follows: (B) Wimmera annual ryegrass (*Lolium rigidum* Gaud. cv. Wimmera), (C) Victorian perennial ryegrass (*Lolium perenne* L. cv. Victorian), (D) Medea perennial ryegrass (*Lolium perenne* L. cv. Medea) and (E) Hybrid phalaris (*Phalaris tuberosa* x *Phalaris arundinacea* cv. Siro 1146).

Large differences between pasture treatments in both pasture and animal productivity occurred in the first year and, to a lesser extent, in the second and third years of set stocking and these could be attributed to differences in botanical composition.

Dominant ryegrass stands in treatments B, C and D following pasture renovation and first year management procedures, proved less productive than the more subterranean clover - dominant pastures in treatments A and E. The data presented show clearly that the availability of the subterranean clover component was the dominant factor in the determination of animal production throughout the experiment.

Medea ryegrass consistently failed to demonstrate any degree of perenniality and Victorian ryegrass plants persisted over no more than three summers.

The influence of increased stocking rate in accelerating the between-year decline in density of Victorian ryegrass plants and the proportion of this species in the pasture, was clearly demonstrated. However, irrespective of stocking rate, the contribution of the three ryegrass species and sown subterranean clovers had declined to negligible proportions by the end of the experiment and this decline was associated with a concurrent increase in the pastures of Yarloop subterranean clover and other annual species, mainly of Mediterranean origin. In particular, the sowing of Mt. Barker and Woogenellup subterranean clovers proved unsuccessful in maintaining a low proportion of Yarloop in the overall legume component of the sward after a period of four to five years. This was highlighted by the level of hard seed reserves of subterranean clover measured in August 1973 in the renovated pasture treatments (mean of c. 210 kg ha⁻¹) of which c. 90 per cent was Yarloop.

The results demonstrated the excellent adaptation of Yarloop subterranean clover to the lateritic podzolic soils of Kangaroo Island: particularly its rapid growth during the winter and early spring and its ability to set copious supplies of seed, irrespective of stocking rate, with potential for germination over a numbers of years. Hence the volunteer annual grass-subterranean clover pastures, typical of this

environment, seemed to be assured a degree of quantitative and qualitative independence of stocking rate.

The results clearly demonstrated an interaction between stocking rate and botanical composition in the determination of animal production following the renovation and early management procedures used in this experiment. However, it was further demonstrated that this was only a short-term effect, because, as the experiment progressed, there was a trend towards like-botanical composition in all pasture treatments.

The efficiency of wool production, in relation to pasture production and rainfall, was dependent on stocking rate and the experiment highlighted the biological and economical inefficiency of low stocking rates.

Most importantly, the experiment showed that provided an adequate legume component existed, change in botanical composition *per se*, resulting from various stocking rates, is no cause for alarm in this environment but rather an indication of the weaknesses of our sown species and the strengths of many of our naturally-occurring volunteer annual pasture species.

INTRODUCTION AND ACKNOWLEDGEMENTS

1. INTRODUCTION

Since European settlement, extensive changes have taken place in the botanical composition and productivity of pastures in southern Australia. The use of superphosphate for top-dressing native grass pastures (Cook 1939) led to a rapid invasion of exotic annual species (including many annual legumes) from the Mediterranean zone. The most notable legume, *Trifolium subterraneum* L. (subterranean clover), provided a substantial lift in soil fertility which further accelerated the invasion of many more of these free-seeding and fast-growing annual grasses and herbs. These invading species have been regarded as weedy and inferior to cultivars of imported temperate grasses (Tiver and Crocker 1951).

Any effort directed towards exploiting the naturalised volunteer pasture species has been, with few exceptions, confined to legumes of undoubted value such as subterranean clover. For example, by 1970, eleven naturally-occurring biotypes (commercial cultivars) of this species were in use in zones with distinct climatic and edaphic features. Some common examples are cv. Yarloop - regions subject to regular flooding: cv. Mt. Barker for long growing seasons: cv. Geraldton for greater speed of seed maturation, etc.

Only recently has the significance of the many attributes of some other common volunteer annuals been recognized (e.g. Carter 1968a and b, Smith 1968b). In particular, their persistence under intensive grazing, and their contribution to pasture and hence animal productivity under diverse management and seasonal influences is fast earning them the reputation of valuable pasture species in many areas of southern Australia.

There appear to have been no reported experiments carried out in the Mediterranean-type environment of southern Australia comparing animal production from pastures based on subterranean clover - volunteer

annual grass and subterranean clover-sown temperate grass.

This report describes an experiment comparing the liveweight response and wool production from wethers grazing at six different stocking rates on subterranean clover - volunteer annual grass pasture with and without the introduction of four temperate grass - subterranean clover mixtures. Particular emphasis has been placed on determining the extent of re-invasion of the renovated pastures by volunteer species, and the persistence of the introduced pasture species, as influenced by stocking rate. An attempt has been made to elucidate some of the inter-relationships of stocking rate and botanical composition as determinants of animal production in the Mediterranean-type environment.

2. ACKNOWLEDGEMENTS

The co-operation of Mr. B. Ninnes and the Kangaroo Island Research Centre staff is gratefully acknowledged and, in particular, field assistance rendered by Messrs. M.R. Lewis, P.G. Telfer and A.D. Murray.

The Australian Wool Corporation substantially financed this project through the Wool Research Trust Fund.

BACKGROUND, AIM, ENVIRONMENT AND SITE

OF EXPERIMENT

3. BACKGROUND TO EXPERIMENTAL PROGRAM

The few relevant experiments reported prior to 1970 showed the extreme lack of evidence for the popular belief that the introduction of perennial grass species into annual pastures in the Mediterranean-type regions of southern Australia, either stabilized pasture composition for an extended period, or resulted in increased animal production. The realization that volunteer annual species, in conjunction with an adequate legume component, may be useful pasture plants was in its infancy.

In addition, little evidence was available on the effect of stocking rate on the rate of invasion of annual volunteer species into sown perennial based pastures or the concomitant decline in perennial grass component. The following experimental work reported in this report has attempted to provide evidence on some of these fundamental questions which are of immediate practical importance to the grazing industries.

A further impetus to the experimental program was the need to solve another important problem which was particularly relevant to Kangaroo Island, South Australia. In the period 1945 to 1970, 100,000 hectares of virgin scrub were cleared for development on Kangaroo Island. The soils are mainly lateritic podzols comprising sandy loams overlying impermeable clay at a depth of from 20 to 60 cm (Northcote and Tucker 1948). The region is typical of the more humid Mediterranean-type environments of southern Australia with precipitation exceeding evaporation during May to August and, due to poor drainage, the soil is generally waterlogged during most of this period. Early experiments carried out on the Kangaroo Island Research Centre, Parndana, demonstrated the outstanding herbage production of Yarloop subterranean clover during the winter (Day 1964) and the use of this cultivar in seed

mixtures was widespread from 1957 to 1963. However, this has resulted in many newly-sown pastures being dominated by Yarloop subterranean clover, and by 1963, the occurrence of a general problem of clover-induced infertility in ewes (Bennetts *et al.* 1946) was recognized on Kangaroo Island, with Yarloop subterranean clover being implicated as the main causal species. The introduction of a successful perennial grass into these pastures could dilute the intake of oestrogenically-potent legume by the grazing animal and perhaps stabilize pasture composition.

4. AIM OF EXPERIMENT

The hypothesis tested in this experiment is specifically that the renovation of existing volunteer annual grass - Yarloop subterranean clover pasture by the tillage, seedbed preparation and sowing of a mixture of the best available grass species and low oestrogenic subterranean clovers results in increased liveweight and wool production from grazing wethers. No attempt has been made to divorce the response due to tillage and/or seedbed preparation from that due to pasture species *per se* in the comparison of renovated and unrenovated treatments.

In essence, the aim of the experiment was to compare the productivity of two systems; one being an unrenovated established pasture comprising subterranean clover - volunteer annual grass (typifying the majority of existing pastures on Kangaroo Island and in many other areas of southern Australia) and the other, a renovated pasture involving tillage, clover seed and grass seed inputs - a new system as defined by the outcome of an established and recommended renovation practice. The experimental comparisons were specifically designed so that the results would be relevant to an established agricultural system; the importance of which had been stressed by Morley and Spedding (1968).

5. ENVIRONMENT

The experiment was located at the Kangaroo Island Research Centre (lat. 35° 48'S., long. 137° 20'E., alt. 153 m.).

Climatic data for the Research Centre are presented in Table 2 and appendix Tables 1-5.

The vegetation of the experimental site was originally a dry sclerophyll scrub dominated by an *E. baxteri* - *E. cosmophylla* association. The main tall shrubs included *Casuarina stricta* (bulloak), *Banksia marginata* (honeysuckle), *Banksia ornata* (broad leaved honeysuckle), *Hakea rostrata* and *Xanthorrhoea tateana* (the endemic yacca). Undershubs included *Daviesia genistifolia*, *D. brevifolia*, *Adenanthos terminalis*, *Lhotskyia glaberrima*, *Phyllota pleurandroides*, *Pultenaea viscidula*, *Logania ovata*, *Spyridium thymifolium*, *Petrophila multisecta*, *Isopogon ceratophyllus*, *Tetratheca halmaturina*, *T. ericifolia* and *Hibbertia stricta* (Northcote and Tucker 1948).

The first morphological description of the soil type was given by Northcote and Tucker (1948) as the Seddon gravelly sandy loam although listed under the incorrect heading of Seddon gravelly loamy sand (refer Carter 1970). The description is as follows:

0-5 cm [†] A0	Scattered surface gravel; occasionally gravel pavement.
A1	Greyish-yellow-brown to grey-brown sandy loam with some organic matter and slight [‡] ironstone gravel.
5-25 cm A2	Yellow-brown to yellowish-brown sandy loam with slight to medium amounts of ironstone gravel.
25-33 cm B1	Light yellow-brown sandy clay loam sometimes with yellow

[†] Depth originally described in inches but converted to centimetres in this text.

[‡] 'Slight' corresponds to 12.5 - 25%, 'light' to 25 - 40% and 'heavy' to more than 70%.

mottlings, with light to heavy amounts of ironstone gravel. (This horizon is frequently poorly developed).

- 33-56 cm B2 Light yellow-brown, brown, and sometimes yellow-grey mottled friable clay. Gravel is very slight.
- 56-102 cm B2 Mottled yellow-brown, brown, yellow-grey and light grey friable clay. Frequently with gravel bands giving medium amounts of ironstone gravel in this horizon.
- 102-152 cm B3 Variously mottled clay but light grey colour prominent, sometimes with ironstone gravel bands.
- 152-203 cm+C Variously mottled clay but red and light grey prominent sometimes with ferruginized rock fragments.

Table 1

SCIENTIFIC NAMES OF ALL SPECIES GIVEN COMMON NAMES IN THE TEXT

<u>Common Name</u>	<u>Scientific Name</u>
Annual ryegrass	<i>Lolium rigidum</i> Gaud.
Annual veldt grass	<i>Ehrharta longiflora</i> Sm.
Barley grass	<i>Hordeum leporinum</i> Link.
Barrel medic	<i>Medicago truncatula</i> Gaertn.
Burr medic	<i>Medicago polymorpha</i> L.
Capeweed	<i>Arctotheca calendula</i> Druce
Cluster clover	<i>Trifolium glomeratum</i> L.
Cocksfoot	<i>Dactylis glomerata</i> L.
Fat hen	<i>Chenopodium album</i> L.
Geranium	<i>Erodium</i> spp.
Goosefoot	<i>Chenopodium</i> spp.
Lucerne	<i>Medicago sativa</i> L.
Meadow foxtail	<i>Alopecurus pratensis</i> L.
Musky crowfoot	<i>Erodium moschatum</i> L'Herit.
Perennial ryegrass	<i>Lolium perenne</i> L.
Phalaris	<i>Phalaris tuberosa</i> L.
Red clover	<i>Trifolium pratense</i> L.
Ripgut brome	<i>Bromus rigidus</i> Roth.
Rough-stalked meadow grass	<i>Poa trivialis</i> L.
Short rotation ryegrass	<i>Lolium perenne</i> L. x <i>Lolium multiflorum</i> Lam.
Silver grass	<i>Vulpia</i> spp.
Siro 1146 hybrid phalaris	<i>Phalaris tuberosa</i> L. x <i>Phalaris arundinacea</i> L.
Soft brome	<i>Bromus mollis</i> L.
Sorrel	<i>Rumex acetosella</i> L.
Subterranean clover	<i>Trifolium subterraneum</i> L.
Tall fescue	<i>Festuca arundinacea</i> Schreb.
Winter grass	<i>Poa annua</i> L.
Yorkshire fog	<i>Holcus lanatus</i> L.

TABLE 2

RAINFALL⁺ RECORDED AT THE KANGAROO ISLAND RESEARCH CENTRE, PARNDANA EAST

YEAR	Monthly Totals (mm)												
	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEPT.	OCT.	NOV.	DEC.	TOTAL
1970	40	3	22	51	50	93	50	112	81	19	27	28	571
1971	6	11	34	138	96	66	62	143	66	37	81	62	792
1972	59	49	6	24	22	63	149	124	18	40	12	6	571
1973	6	8	71	42	62	108	90	140	72	56	18	43	699
1974	50	8	25	60	109	66	161	81	72	93	27	9	762
20 year Ave.	16	25	27	53	78	87	104	88	62	44	28	26	638

⁺ Other climatic data are given in appendix tables 1 to 5.

6. SITE OF EXPERIMENT

The experimental site was cleared of native vegetation in 1950 by the South Australian Lands Development Executive and until 1962 had been used for experiments evaluating response to various rates and frequency of superphosphate and copper fertilizer applications. From 1963 to 1969 inclusive, the entire site was managed as a single paddock and received uniform applications of fertilizer. Experiments carried out on the Research Centre on the same soil type indicated that the possible optimum maintenance superphosphate requirement for pasture growth was between 250 and 375 kg ha⁻¹ year⁻¹ (Carter 1970). The maximum and minimum mean rates of fertilizer applied to any portion of the site prior to 1970 were 243 and 154 kg ha⁻¹ year⁻¹ respectively of superphosphate and 0.9 and 0.3 kg ha⁻¹ year⁻¹ respectively of copper as copper sulphate. Hence the mean annual application of superphosphate was below the estimated optimum annual maintenance requirements for pasture production. Although the phosphate sorption capacity is reduced by application of phosphatic fertilizers (Kanwar 1956, Kurtz and Quirk 1965) this reduction is insignificant compared to the extremely large phosphate retention capacity of this soil i.e. 15,400 p.p.m. for virgin 0-10 cm soil and 11,250 p.p.m. for partially-developed 0-10 cm soil (Kanwar 1956). However, to minimize possible variation in pasture production due to previous variation in fertilizer application on the trial site, a blanket dressing of 348 kg ha⁻¹ of superphosphate and .94kg ha⁻¹ of copper was applied in 1970, during the establishment of the experiment.

A pasture mixture of 4.5 kg ha⁻¹ of Bacchus Marsh subterranean clover and a trace of Colac perennial ryegrass and GB 81 phalaris had been sown on the newly-cleared land in 1951 but these species were

scarce on the site in 1970. In 1960 the site was worked up in the spring, stumps ripped and the soil levelled and in 1961 a mixture comprising perennial ryegrass and Yarloop and Mt. Barker subterranean clovers sown. By 1969 the pasture was dominated by Yarloop subterranean clover and barley grass.

Other species present were Mt. Barker, Dwalganup (although never sowed on the site) and Bacchus Marsh subterranean clovers, silver grass, soft brome, riggut brome, geranium and capeweed.

EXPERIMENTAL METHODS

7. EXPERIMENTAL METHODS

7.1 Site preparation

An area of 11.6 hectares of land was prepared in 1970, with the exception of certain areas allocated to an unrenovated pasture treatment (see 7.2.1), as follows:

Dried pasture residue remaining from the 1969 growing season was burnt in March, ploughed in early April, cultivated and harrowed in mid-April and finally rolled with a cultipacker in late April.

7.2 Treatments

7.2.1 Pasture treatments

Five pasture treatments were incorporated in the design as follows:

Treatment A: The unrenovated, volunteer annual grass - subterranean clover pasture that was present on the experimental site as described in 6. The paddocks comprising this treatment were untouched in any way during the renovation of the remaining four treatments and for the duration of the experiment (except for the topdressing of fertilizer).
Treatments B, C, D and E: Tillage, seed bed preparation and sowing of respective treatment grasses together with Mt. Barker and Woogenellup subterranean clovers. Treatment grasses were as follows:

B: Wimmera annual ryegrass

C: Victorian perennial ryegrass

D: Medea perennial ryegrass

E: Siro 1146 hybrid phalaris

The pasture treatments will hereafter be referred to as A, B, C, D and E respectively.

Wimmera and Victorian ryegrasses have been and are the most commonly sown annual and perennial grasses respectively on Kangaroo Island.

Cutting experiments carried out on the Kangaroo Island Research Centre had shown Victorian ryegrass to be the most productive perennial pasture species in this environment (Crawford 1956). Tall fescues and cocksfoots had generally produced only small quantities of dry matter compared to the ryegrasses (Crawford 1956) and great difficulty had been encountered in adequately establishing varieties of phalaris on the ironstone soils.

Medea ryegrass was a recently released cultivar developed at the Waite Institute (Silsbury 1961) by selection of plants collected near Medea in Algeria (Neal-Smith 1955). This cultivar was reputed to be characterized by a high degree of summer dormancy and persistence (Barnard 1972) and would have potential for widespread use on Kangaroo Island if its reputation for persistence could be substantiated under grazing on this soil type. Trials in Victoria had also shown Medea to produce more dry matter in late autumn, winter and early spring than Victorian ryegrass but that it was less productive during the rest of the season (Cade 1969).

Siro 1146 hybrid phalaris was developed by the CSIRO by the hybridization of *Phalaris tuberosa* and the hexaploid race of *Phalaris arundinacea*. Details concerning the hybridization have been reported in several publications (McWilliam 1962, McWilliam and Neal-Smith 1962, Anon 1962). The more recent results concerning the characteristics and performance of this hybrid have been described (McWilliam *et al.* 1965, Myers and Squires 1968b). Prior to 1970 this pasture species had received little testing under grazing but in the few trials that were then in progress the hybrid had shown an ability to establish readily, to combine well with a variety of legumes and to persist under heavy stocking and through extended dry periods. Although it generally appeared that maximum benefit from the use of the hybrid would be obtained from

regions where summer rainfall incidence was high or where irrigation was available, reports of superior ease of establishment and persistence under waterlogged conditions indicated that this species may have been useful in the Kangaroo Island environment.

7.2.2 Stocking rates

Information derived from previous experiments carried out on the Kangaroo Island Research Centre (Carter 1965, Carter and Day 1970) indicated that large-framed, strong-wool type, South Australian Merino wethers could be continuously grazed without supplementary feeding at a stocking rate of 13 sheep ha⁻¹ without undue stress on the sheep. This was based on the pasture containing a major component of subterranean clover and supplied with an annual maintenance dressing of superphosphate at c. 190 kg ha⁻¹.


Six stocking rates chosen for the experiment described in this report were 10, 11, 13, 14, 15 and 17 sheep ha⁻¹, giving a range, incorporating an average and maximum, slightly higher than the respective 13 and 16.5 ha⁻¹ used in the grazing experiment of Carter and Day (1970).

7.2.3 Layout and allocation of treatments

Pasture treatments and stocking rates were allocated at random to the experimental site which comprised fifteen elongate paddocks on each side of a single raceway. The length of each paddock was constant and the width varied to give the required areas for the allocated stocking rates. Seed bed preparation and sowing of pastures was completed prior to the erection of fences and watering facilities. Once set stocked, each paddock was continuously grazed by five sheep thus utilizing 150 experimental animals. There was no replication of treatments:

Plan of experimental paddocks

Pasture treatments and stocking rates were allocated at random to the experimental site. The length of each paddock was constant and the width varied to give the required areas for the allocated stocking rates. Each pasture treatment was unreplicated at comparative stocking rates.



Treatments

- A - Unrenovated
- B - Wimmera ARG
- C - Victorian PRG
- D - Medea PRG
- E - Hybrid Phalaris

Postscripts for treatments indicate stocking rates in wethers hectare⁻¹

i.e. the 30 paddocks comprised Pastures (5) x Stocking Rates (6) = 30.

7.3 Pasture establishment

Pastures were sown on May 11, 12 and 13, 1970, as grass-subterranean clover mixtures through the seeds box of a 12 row, E series, Shearer combine. All seed, except Wimmera ryegrass, was certified with a potential germination of 90%[†].

The experimental site had not previously been sown to Woogenellup subterranean clover: therefore, because of its specific rhizobial requirements, all Woogenellup seed was inoculated with commercial inoculant prior to sowing. Sowing rates used were approximately those currently recommended by the South Australian Department of Agriculture and adjusted to a 90% germination. Actual rates were:

- Wimmera ryegrass (Treat. B) 19.4 kg ha⁻¹.
- Victorian ryegrass (Treat. C) 10 kg ha⁻¹.
- Medea ryegrass (Treat. D) 9.7 kg ha⁻¹.
- Siro 1146 hybrid phalaris (Treat. E) 4.3 kg ha⁻¹.

Woogenellup and Mt. Barker subterranean clovers were sown in the ratio 3:2 at a combined sowing rate of 10.4 kg ha⁻¹.

7.4 Fertilizer application

All paddocks received 348 kg ha⁻¹ of normal single superphosphate (9.6% P, 22% P₂O₅) during pasture establishment in May 1970, and applied as follows:

In treatments B, C, D and E 140 kg ha⁻¹ was drilled in with the seed and in treatment A 140 kg ha⁻¹ was topdressed onto the pasture. All treatments were topdressed again with 208 kg ha⁻¹ of superphosphate

[†] Tested by the South Australian Department of Agriculture and Fisheries' seed testing laboratory.

including 0.94 kg ha^{-1} of copper and 30 g ha^{-1} of molybdenum.

7.5 Selection and allocation of experimental animals

In October 1970, a line of 1000 wether hoggets (South Australian strong wool Merinos - Collinsville blood) from Pitlochry Cortina, Cantara, South Australia, were visually examined for any obvious foot, mouth, pizzle or wool defects and 370 culled from the main flock. The remaining 630 sheep were consecutively numbered and shorn. Total greasy fleece weight (weighed to 0.05 kg), including locks, skirtings and pieces but excluding belly wool and dags, was recorded for each sheep. The wethers were then weighed (to the nearest 0.5 kg) off-shears and 250 which conformed to the criteria, $40.9 \text{ kg} < \text{live weight} < 52.2 \text{ kg}$ and $5.3 \text{ kg} < \text{greasy fleece weight} < 6.8 \text{ kg}$, were purchased.

Following transport to the Kangaroo Island Research Centre the wethers were given a cobalt bullet, drenched, tagged with large, easily-read cattle tags and grazed as a single flock until placed on their allocated paddocks on December 22, 1970. The allocation of sheep to paddocks was performed by graphically plotting the fleece weight against live weight for each sheep (as determined in October, 1970) and stratifying the 250 wethers into groups. Allocation to both pasture treatments and stocking rates was made from these groupings at random. Each of the 30 paddocks comprising the experiment was allocated five sheep with similar mean live weight and mean greasy fleece weight at the start of the experiment. The 100 wethers remaining were run as one flock of spares in laneways and spare paddocks around the experimental site. The wethers were two years old at the time of placement in their paddocks and thereafter they grazed their allotted paddocks continuously until the end of the experiment in January 1975.

7.6 Grazing management of pastures in 1970

Management of the pastures in the first year prior to set stocking with the experimental animals was designed to fulfil the following criteria.

a) Ensure that approximately equivalent amounts of pasture were available on all paddocks prior to the allocation of experimental animals to their paddocks in December 1970.

b) Ensure that the treatment grasses were dominant in 1971 (i.e. allow maximum seed set of treatment grasses in the spring of 1970) without undue suppression of the subterranean clover component.

c) Graze treatment A paddocks at least once in keeping with normal commercial practice.

Treatment A paddocks were uniformly grazed for six weeks (June 18 to July 29) by a large flock of wethers divided into equivalent numbers per unit area for each paddock. Treatment B paddocks were similarly grazed during the early spring (September 17 to October 7) as pasture production was far in excess of the other renovated treatments and severe suppression of the subterranean clover component was occurring. The allowance of an adequate growth period following the grazing of treatment B was considered essential to allow adequate seed set of the Wimmera ryegrass. Treatments C, D and E remained ungrazed until set stocked with the experimental animals in December 1970.

7.7 Routine management

7.7.1 Pastures

Fertilizer application: From 1971 to 1974 inclusive, all paddocks were topdressed with 200 kg ha^{-1} of single superphosphate prior to the opening rains in each year.

Insect control: All paddocks were misted with 350 g ha^{-1} (active ingredient) of 'Imidine' five weeks after the opening rains in each year to control red-legged-earth mite (*Halotydeus destructor*). In some years a second misting was performed to control later hatchings.

7.7.2 Animals

Flystrike: The wethers were crutched once during the spring of each year and strategically jetted (common treatment to all sheep) with 'Diazinon' in the experimental paddocks as required.

Worms: All experimental animals were drenched in the autumn of each year with 'Thiobendazole' to control round worms. Additional drenching was carried out as required.

Lice: Dipping was carried out off-shears for control of lice (*Damalina ovis*) and ked (*Melophagus ovinus*) prior to selection of the experimental animals and again in January 1973.

Pizzle rot: During the experimental period a number of animals were lanced to release accumulated fluids.

Supplementary feeding: Wethers grazing in treatments B, C and D at the highest stocking rate (17 ha^{-1}) were fed $0.5 \text{ kg of oats sheep}^{-1} \text{ day}^{-1}$ for one week only, commencing 2nd July 1972, to prevent death from undernutrition. No hand feeding or fodder conservation was carried out at any other time during the experiment.

Replacement of animals: Experimental animals that died in their paddocks were replaced within a few days (usually within 24 hours) from the reserve flock grazing near the experimental site. The replacement sheep were matched as closely as possible on the basis of live weight and

wool cutting ability as determined in the original weighing and shearing (October 1970).

Live weights of replacement sheep were not used in the group data means for at least 12 weeks. Fleece weights and measurements of fleece characteristics of replacement animals were not used in the data analysis in the year of replacement.

Causes of death of experimental wethers other than undernutrition included pizzle rot, obesity and flystrike and are listed in appendix Table 29.

7.8 Data collection procedures

7.8.1 Pasture data 1970

Established plants of the sown pasture species and Yarloop subterranean clover were counted in each of ten randomly-thrown, 0.04 m^2 quadrats on June 17, 1970. Pasture production in the sown treatments was measured over two periods i.e. May 11 to September 17 and September 17 to November 25 using the techniques described for 'open' cuts in 7.8.2. In treatment B both 'open and closed' cuts were taken because of the presence of grazing sheep as described in 7.6.

Available pasture was measured on all paddocks on November 11 to coincide with the end of the growing season.

7.8.2 Pasture data 1971-1974

The 'open and closed quadrat' method of McIntyre (1946) was used to estimate dry matter availability, seasonal and total dry matter production, and crude utilization (amount of pasture dry matter eaten or otherwise disappearing - refer Carter and Day 1970).

Samples were cut from six areas protected by 'Weldmesh' cages

(closed quadrats) c. 1.4 m square. Within each of these cages a 0.5 m² quadrat was cut at ground level with a mobile, motorized sheep-shearing hand piece with 'wide' combs and cutters. A corresponding sample was cut from without, but in close proximity to, each cage. Thus the number, size and shape of the 'closed' cuts from within the cages were the same as for the 'open' quadrats. Following cutting of the protected quadrat site, the cage was placed at random on a new un-cut site in the paddock. No pasture measurements were made during the characteristic 4-5 month period of summer drought and the protective cages were shifted to a new site just before the opening rains in each season. Three samplings were made during each growing season to approximately coincide with the end of the three calendar seasons, autumn, winter and spring. The one exception being in 1971 when an extra sampling was made in July. On most occasions, individual green weights were recorded for each of the six 'open' and six 'closed' samples. These samples were then bulked into 'open' and 'closed' lots and a representative subsample was taken from each lot. The size of the subsample was usually 10% of the total green herbage within each lot but varied depending on size and growth stage of the pasture plants. Strict mixing and subsampling procedures were followed to ensure representative subsamples.

Each subsample was hand-separated, while fresh, into botanical components which were then dried separately to constant weight in a forced air-drought oven at 85°C.

The components were as follows:

- (1) Sown grass
- (2) Subterranean clover
- (3) Broad-leaved volunteers
- and (4) Narrow-leaved volunteers.

Plant density: Ten fixed sampling sites marked with yellow-painted jarrah pegs were selected at random in each paddock at the start of the

experiment. In June of each year all perennial[†] plants and subterranean clover plants within a one m⁻² quadrat, placed at each site, were counted.

7.8.3 Subterranean clover seed

Measurement of subterranean clover seed present in, or on, the soil was made on four occasions during the experiment, i.e. August 1973 and 1974 and January 1974 and 1975. The difference in quantity of seed recorded between the spring and following summer sampling was used as a crude estimation of seed produced during the season. No attempt was made to measure loss of seed due to trampling, eating, or any other cause, during the intervening period. The sampling during August also estimated the hard seed reserves present at that time.

At the August samplings 30 round cores of 10 cm diameter and 5 cm depth were taken at random within three transects across each paddock. The 10 cores from each paddock transect were bulked to give three lots per paddock which were then dried at 50°C, broken up and sieved through a 0.5 mm sieve to reduce the sample bulk by separation of fine soil particles. The sample was then thoroughly stirred in 'Genklene' solution (trichloroethane SG 1.314) to separate the coarse sand, gravel and stones from the floating sample mass (Carter, Challis and Ridgway 1977).

The sample was then air dried, threshed in a belt thrasher, hand-sorted into white seed (Yarloop) and black seed (non-Yarloop) and each lot weighed to 0.001 g.

At the January samplings, above and below-ground components were sampled separately. The above-ground sampling was performed by cutting six 0.5 m² quadrats at ground level in each paddock in a similar manner to that described for pasture availability during the growing season. The soil surface was then brushed with a fibre brush and all loose soil and dry herbage collected, dried at 50°C for 24 hours and hand-picked to

[†] plants that had survived at least one summer.

remove faeces, sticks and large stones. Following the removal of the above-ground material, the soil was carefully hoed to a depth of 5 cm, removed and bulked into three sample lots per paddock. Final treatment of the above and below-ground fractions was similar to that used for the August samples.

7.8.4 Sheep live weights

All sheep (experimental and spares) were weighed at approximately six-weekly intervals. Occasionally, additional weighings were made when animals were in stress periods. The sheep were weighed in the raceway, at the entrance to each paddock, using mobile scales. Liveweight data were recorded to the nearest 0.1 kg. Sheep were weighed throughout the day and no special measures were taken to ensure sheep were of similar gut-fill.

7.8.5 Annual wool production

All experimental sheep were removed from their paddocks on the day of, or one day prior to, shearing and were driven as a single flock to the shearing shed (c. 1 km). The sheep were transported back to their paddocks, the same day or early the next day following shearing, in their respective experimental groups. All sheep were combined into one flock in the shearing shed and randomly selected by the shearers. The same procedure was repeated in the spring of each year when the sheep were crutched and only on these two occasions, during any one year, were the experimental sheep removed from their paddocks.

Total greasy fleece weight in all data presented is the summation of the individual weights of dyeband sample (refer 7.8.7), midside sample (refer 7.8.6) locks, belly and skirted fleece of each locks and fleece of each sheep were weighed separat

the exception of the January 1972[†] shearing when the total weight only was recorded).

7.8.6 Fleece characteristics

A sample of wool was taken from the midside of each fleece as each sheep was shorn. This sample was then weighed, placed in an air-tight plastic bag with an identification card, and sent to the Gordon Institute of Technology, Geelong, Victoria for analysis. The following fleece measurements were made on each midside sample: a) Yield at 16% regain b) mean staple length c) mean number of crimps per unit length of fibre (crimp frequency) and d) mean fibre diameter.

7.8.7 Seasonal wool production

On most occasions of liveweight measurement, a dyeband (10-15 cm long) was applied dorso-ventrally on each sheep in the approximate position of the last rib, according to the technique of Short and Chapman (1965).

Each year, one day prior to annual shearing, the dyeband staples were carefully removed by clipping the wool as close to the skin as possible using 'Oster' small animal clippers. Application of dyebands and removal of dyeband staples were performed in a sheep-holding cage. This unit contained a rotatable side to which the animal was strapped and held on its side at a convenient working height. The width of staple removed was from c. 2-3 cm of skin. Each harvested, dyebanded staple was weighed, had a length of tissue paper placed along its length, rolled and stored in a bag.

All dyeband staples for all years were measured as follows:
The staples were opened so that an even spread of dyebands was obtained on both sides. Five measurements were made on each half of the sample

[†] shearing was in January each year (refer appendix Tables 17-20 for dates).

(taking care to measure staples of good length, i.e. away from the belly and back extremities). The staple was placed in a natural (unstretched) position and the measurements were made by zeroing a ruler on the base of the staple, and then reading off the lengths at the proximal end of each dyeband. The length between dyebands, used in conjunction with the total clean fleece weight, allowed estimates to be made of periodic clean wool growth during each year.

7.9 Data analysis

In all statistical analyses, the mean measurement for each paddock has been used as the basic experimental unit (e.g. mean live weight of the five sheep on each weighing occasion, or mean availability of pasture in each paddock on each sampling occasion). Regressions of each unit of measurement on stocking rate on each sampling occasion were performed and the relationship tested for linearity. On nearly all occasions, higher order polynomial relationships could not be justified statistically due to insufficient levels of the x variate (stocking rate), even though they may have been biologically possible or probable. Tests of homogeneity of regression coefficients (homogeneity of the slopes of the regression of each unit of measurement, for each pasture treatment, on stocking rate) were performed to indicate any significant interaction between pasture treatment and stocking rate. The mean regression coefficient for the five regression relationships (five pasture treatments on stocking rate) has also been calculated for each unit of measurement.

Where the heterogeneity of regression coefficients was not statistically significant ($P > 0.05$) a test of homogeneity of adjusted treatment means (adjusted to $x = \bar{x}$ where \bar{x} = mean of all the x-values of K treatment samples⁺) was performed to indicate whether the pasture

⁺ as the same stocking rates were used over each pasture treatment
 $\bar{x} = \bar{x}$ (mean of stocking rates 10, 11, 13, 14, 15 and 17 ha⁻¹).

treatment means (averaged over stocking rates) were statistically different and the least significant difference (LSD) between these means calculated. Angular transformation of proportional data (percent) was performed as required.

RESULTS

8. RESULTS

8.1 Pasture establishment

With the exception of the hybrid phalaris, the sown grasses and legumes established satisfactorily (Table 3), in particular, the density of perennial ryegrass plants (treatments C and D) were obviously higher than the density that could be maintained as perennial plants in the following years. The density of established hybrid phalaris plants was low, although superior to previously obtained densities of phalaris (cv. Australian) sown at similar rates on this soil type on Kangaroo Island. The density of hybrid plants was considered sufficient to allow this species to make a worthwhile contribution to pasture production in the later years of the experiment, provided that the individual plants expanded in diameter, as is characteristic of this species.

There was no significant ($P > 0.05$) difference between the density of established sown grass plants within each sown pasture treatment over the range of stocking rates, or in the case of subterranean clover plants, over all renovated paddocks. However, the mean density of Yarloop subterranean clover plants in treatment A was $580 \text{ plants m}^{-2}$ compared to the combined 85 plants m^{-2} of Mt. Barker and Woogenellup and 25 plants m^{-2} of Yarloop subterranean clovers established in the renovated paddocks.

A visual inspection of the pastures during the spring showed that, within the narrow-leaved volunteer component, barley grass and ripgut brome were the dominant species in treatments A and E respectively.

8.2 Pasture production 1970

The superior pasture production in treatment B during the autumn and winter is shown in Table 4. There was less difference between pasture treatments in pasture production during the spring. A different seasonal

growth pattern for the two perennial ryegrass species (treatments C and D) is also evident in Table 4.

Some variation in levels of available pasture was present at the end of the 1970 growing season (Table 4) because of the difficulty of achieving even grazing. Treatment A contained the lowest, and treatment D the highest, amounts of available pasture at the end of the 1970 growing season.

8.3 Botanical composition 1971-1974

8.3.1 Sown grasses

The renovation of the original pasture by the sowing of Wimmera ryegrass (treatment B), Victorian ryegrass (treatment C) and Medea ryegrass (treatment D), together with Mt. Barker and Woogenellup subterranean clovers resulted in ryegrass-dominant pastures in the autumn of 1971 (Table 5).

The majority of Victorian ryegrass plants died over the 1972/73 summer (Table 6) and although some self-sowing of this species occurred in the autumn of 1973, its contribution was negligible by the end of the 1973 season (Table 5). The Medea ryegrass plants died during each summer of the experiment except for a few plants which survived the 1973/74 summer at the lower stocking rates (Table 6).

In the first three years of set stocking there was a within-year decline in the ryegrass component in all three ryegrass treatments during the autumn and winter and in some years the decline continued during the spring.

An effect of stocking rate on the between-year decline in the proportion of Wimmera and Victorian ryegrasses was evident over the first two summers following set stocking (1971/72 and 1972/73) Figure 1.

Table 3

PURITY, VIABILITY AND ESTABLISHMENT OF SOWN PASTURE SEED 1970

Species	Purity %	Germ ⁿ 14 days %	Viable Sowing Rate (kg ha ⁻¹)	Potential Establish. (plant m ⁻²)	Actual ⁺ Establish. (plant m ⁻²)	Establish. ⁺ %
Wimmera A.R.G.	-		19.4	1000	290	29.0
Victorian P.R.G.	99.8	90	10.0	600	190	31.6
Medea P.R.G.	99.5	90	10.0	600	200	33.3
Hybrid Phalaris	99.0	88	4.3	375	56	15
Mt. Barker	99.3	90	6.2	((110	((85	((77
Woogenellup	99.6	92	4.2	(((

⁺ established plants on June 17, 1970.

Table 4

PASTURE GROWTH FOR TWO PERIODS IN 1970 AND AVAILABLE PASTURE AT THE
END OF THE 1970 GROWING SEASON[†]

Growth Period	Pasture treatment means					LSD 5%
	A	B	C	D	E ₁	
	Pasture growth (kg ha ⁻¹ day ⁻¹)					
11/5-17/9	-	44	21	33	18	3.2
17/9-25/11	-	68	65	51	59	10.7
Date	Pasture availability (kg ha ⁻¹)					
25/11	4575	6230	7210	7750	6360	570

[†] Pasture treatments A and B received 1600 and 3600 grazing days ha⁻¹ respectively





Figure 1. Effect of stocking rate and duration of continuous grazing on the botanical composition measured in each treatment on three occasions in each year of the experiment.



+ = mean of 14, 15 and 17 wethers ha⁻¹

* = mean of 10, 11 and 13 wethers ha⁻¹

Table 5

PROPORTION OF SOWN GRASS IN THE PASTURE ON TWELVE OCCASIONS DURING THE EXPERIMENT

(% Dry Weight)

Date	Pasture Treatments				Hetero. Adj. Treat. Means	LSD 5%	Sig. Mean Reg. on SR.	Hetero Reg. Coeff.
	B	C	D	E				
2/6/71	97	91	97	15	***	5.6	NS	NS
8/9	90	83	97	8	***	9.9	NS	NS
24/11	*+ 51	45	73	6	***	14.5	NS	NS
27/6/72	56	** 82	83	22	***	18.4	*	NS
26/9	13	51	37	8	***	9.6	NS	NS
16/11	15	26	34	8	***	9.6	NS	NS
18/6/73	* 35	24	37	9	-	-	**	*
17/9	18	7	9	7	*	9.4	*	NS
26/11	19	9	* 23	12	-	-	*	*
10/6/74	4	3	4	12	NS	8.6	NS	NS
2/9	2	3	5	7	NS	3.8	*	NS
18/11	7	5	* 12	21	-	-	**	**

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

+ = Significance of within-treatment regressions on stocking rate

Table 6

DENSITY OF PERENNIAL⁺ PLANTS PRESENT IN JUNE OF EACH YEAR

(Mean of 10 fixed locations per paddock)

(plants m⁻²)

Year	Stocking Rate						Sig.† Reg.
	10	11	13	14	15	17	
	Victorian perennial ryegrass						
1971	48	47	51	54	47	49	NS
1972	47	38	40	30	25	21	**
1973	5	2	1	0	0	0	*
1974	10	12	8	6	4	1	**
	Medea perennial ryegrass						
1971	0	0	0	0	0	0	-
1972	0	0	0	0	0	0	-
1973	0	0	0	0	0	0	-
1974	15	12	7	6	5	5	**
	Hybrid phalaris						
1971	25	29	28	28	34	30	NS
1972	18	23	20	19	26	20	NS
1973	17	22	20	18	25	19	NS
1974	17	21	19	18	26	18	NS

+ = Plants which have survived at least one summer

† = Significance of regression of plant number on stocking rate.

NS = P > 0.05, * = P < 0.05, ** = P < 0.01

In each case, higher stocking rates were associated with a lower sown grass component in the autumn of the following year.

In the Victorian ryegrass treatment the decline in proportion of ryegrass was usually associated with a between-year decline in the number of perennial plants that survived each summer and increased stocking rate enhanced this between-year decline in density of Victorian ryegrass plants (Table 6).

However, irrespective of stocking rate, the contribution of the ryegrass in treatments B, C and D had declined to almost negligible proportions by the 1974 season (Table 5).

In contrast to the three sown ryegrass treatments, the renovation of the original pasture by the sowing of the hybrid phalaris (treatment E) resulted in only a low proportion of this species component being present in the pasture in the autumn of 1971 (Table 5). The size of the hybrid phalaris component varied little over the experimental period and, although in most years there was a larger component at the lower stocking rates (Figure 1), on no occasion was the regression of the proportion of hybrid phalaris in the pasture, or density of perennial plants (Table 6) on stocking rate, significant ($P > 0.05$). Following an initial between-year decline from 1971 to 1972 the mean number of hybrid phalaris plants remained constant, although it was noticeable from visual observation that the crown circumference of the plants had increased over the experimental period, particularly at the lower stocking rates.

8.3.2 Subterranean clover

The proportion of subterranean clover in the pasture in 1971 was highest in treatment A and lowest in treatment D and reflected the management of the pastures in 1970. The vigorous growth of the sown ryegrasses in 1970 resulted in suppression of the clover growth (and

probably seed production in the spring). The less vigorous growth of hybrid phalaris in treatment E did not result in the suppression of clover growth in 1970, and allowed adequate seed set of the introduced subterranean clover varieties (Mt. Barker and Woogenellup). Consequently, in 1971 the density of clover plants in treatment E, as also in treatment A, was significantly ($P < 0.001$) higher than in the three ryegrass treatments (Table 7). However, all pasture treatments contained a substantial clover component by the spring of 1972 (Figure 1) associated with a rapid between-year increase in the density of clover plants in treatments B, C and D from 1971 to 1972 (Table 7).

The proportion of Yarloop in the total subterranean clover component averaged 27 per cent for the four renovated treatments (B, C, D and E) in June 1971 but the rapid invasion of Yarloop into these treatments (irrespective of stocking rate) resulted in this proportion being increased to 76 per cent by June 1974 (Table 8).

In treatment A the early physiological maturation of Yarloop resulted in a within-year decline in the clover component during the late spring following a peak contribution during the winter in all years (Figure 1). In contrast there was a general pattern of an increasing within-year subterranean clover component throughout the growing season in the renovated treatments, particularly in treatment E (Figure 1). In 1973 there was a marked decline in the clover component during the spring in all treatments (Figure 1) due to a severe attack of clover scorch caused by the fungus *Kabatiella caulivora* (Walker 1956, Beale 1972).

Only on one occasion (September 26, 1972) during the experiment was a significant ($P < 0.05$) interaction between pasture treatment and stocking rate in the proportion of clover present in the pasture recorded (Tables 9 and 10). On this occasion a significant ($P < 0.05$) increase in

proportion of subterranean clover with increased stocking rate was recorded in treatment A but not in any of the other pasture treatments.

On November 24, 1971, November 26, 1973 and on all sampling occasions in 1974 the mean proportion of clover in the pasture increased significantly with increased stocking rate (Tables 9 and 10). The mean regression coefficient (mean proportion of clover over all pasture treatments on stocking rate) differed significantly from zero (Tables 9 and 10). However, with the exception of treatment D, no influence of stocking rate on the density of clover plants counted in June of each year was evident (Table 7). In treatment D, a decrease in the proportion of clover with increased stocking rate was evident on four sampling occasions from September 26, 1972 to September 17, 1973 inclusive although the regression was only significant ($P < 0.05$) on one occasion (June 18, 1973). This relationship was associated with a significant decline in density of clover plants with increased stocking rate in treatment D in 1972 and 1973 (Table 7). The data in Table 7 also emphasizes the lower mean density of clover plants in treatment D in 1971, 1972 and 1973 compared to the other pasture treatments.

8.3.3 Volunteer annual species

Treatment E was rapidly invaded by volunteer annual species (particularly ripgut brome) during 1970, the year of pasture establishment, due to the poor establishment and seedling vigour of the hybrid phalaris. In 1971, the broad-leaved component, comprising mainly capeweed and geranium, was greatest in this treatment, particularly at the lower stocking rates (Figure 1, Table 11). However, in both treatments A and E the narrow-leaved component, comprising mainly barley grass and ripgut brome respectively, was the dominant pasture component in 1971 (Table 13). The broad-leaved component increased from one year to the next in all

treatments and by 1974 was making a significant contribution to the botanical composition (Figure 1, Tables 11 and 12). Particularly evident was the between-year increase in the broad-leaved component from 1972 to 1973 in treatment C (Figure 1, Tables 11 and 12) following the death of the Victorian ryegrass plants over the 1972/73 summer.

The mean (of treatments) broad-leaved component was significantly ($P < 0.01$) larger at the high stocking rates on the first sampling occasion in 1973 (June 18, 1973) but by the final sampling occasion in the spring of the same year (November 26, 1973) the relationship was reversed with a significantly ($P < 0.01$) smaller component at the high stocking rates (Figure 1, Table 12). However, no effect of stocking rate on the proportion of broad-leaved volunteers in the pasture was evident in 1974.

On three occasions in 1971 and 1972, significant interactions between stocking rate and pasture treatment in the size of the narrow-leaved volunteer component were recorded (Table 13). This interaction was mainly due to a decline in the narrow-leaved component in treatment A and increase in treatment E with increased stocking rate resulting in no record of any significant ($P > 0.05$) mean (over all treatments) influence of stocking rate on the size of the narrow-leaved component. In 1973 and 1974, instances of stocking rate affecting the size of the mean narrow-leaved component were recorded (Figure 1, Table 14); however the relationship was positive in 1973 and negative in 1974 (Table 14). In treatment E in 1974, as in 1971 and 1972, the size of the narrow-leaved component increased with increased stocking rate (this component tended to decrease with increased stocking rate in the other pasture treatments) resulting in a significant Pasture treatment x Stocking rate interaction being recorded on two occasions (Table 14).

Table 7

DENSITY OF SUBTERRANEAN CLOVER PLANTS PRESENT IN JUNE OF EACH YEAR

(plants m^{-2})

Year	Pasture treatments means					LSD 5%
	A	B	C	D	E	
1971	534	92	81	8	362	64
1972	438	411	288	**+ 172	318	87
1973	356	251	284	* 195	373	51
1974	233	243	267	242	255	46

+ = Significance of within-treatment regressions on stocking rate.

* = $P < 0.05$, ** = $P < 0.01$

Table 8

PROPORTION OF YARLOOP SUBTERRANEAN CLOVER IN THE TOTAL

CLOVER COMPONENT PRESENT IN THE PASTURES IN

1971 AND 1974

(% dry weight)

Date	Pasture treatment means					LSD 5%
	A	B	C	D	E	
2/6/71	100	30	20	37	22	6.4
10/6/74	100	81	70	87	66	10.4

Table 9

PROPORTION OF SUBTERRANEAN CLOVER IN THE PASTURE ON SIX OCCASIONS IN 1971-1972
(% dry weight)

Date	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)						Grand Mean	Hetero. Adj't. treat. mean	LSD 5% treat. mean	Sig. mean	Hetero. Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15						17
2/6/71	18	2	1	0	8	2	5	8	7	6	6	6	***	5.2	NS	NS
8/9	53	10	7	1	23	16	18	21	23	16	18	19	***	7.9	NS	NS
24/11	27	41	27	7	44	19	27	32	25	36	36	29	***	12.2	**	NS
27/6/72	24	35	9	10	18	10	21	16	20	22	26	19	**	13.6	NS	NS
26/9	33 ⁺	70	38	37	29	38	38	46	40	44	42	41	-	-	NS	*
16/11	20	61	51	32	46	33	50	51	32	49	38	42	***	15.8	NS	NS

⁺ = Significance of within-treatment regressions on stocking rate.

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Table 10

PROPORTION OF SUBTERRANEAN CLOVER IN THE PASTURE ON SIX OCCASIONS IN 1973-1974

Date	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)						Grand Hetero. mean	Hetero. Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15	17			Hetero. Adj. Treat. means
18/6/73	44	33	36	† 35	40	41	35	38	39	40	31	37	11.7	NS
17/9	81	59	69	75	67	70	60	67	78	76	71	70	18.2	NS
26/11	7	13	† 22	14	20	10	9	11	23	15	22	15	13.2	*
10/6/74	16*	17	10	13	10	4	2	12	16	25	19	13	12.4	**
2/9	29*	22	27	29	30	14	7	28	38	35	42	27	15.4	***
18/11	18*	25	36	34	42	21	14	26	35	48	42	31	13.9	***

+ = Significance of within-treatment regressions on stocking rate.

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

† = negative correlation

Table 11

PROPORTION OF BROAD-LEAVED VOLUNTEERS IN THE PASTURE ON SIX OCCASIONS IN 1971-1972

Date	(% dry weight)																	Grand Hetero. mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.			
	Pasture treatment means						Stocking rate (sheep ha ⁻¹)					Grand Hetero. mean													
	A	B	C	D	E	10	11	13	14	15	17														
2/6/71	4	0	2	0	**+	6	8	3	3	1	1	3.6	-	-	**	**	**								
8/9	2	1	4	0	15	10	2	7	4	2	2	4.5	*	9.8	NS	NS	NS								
24/11	0	0	0	0	2	2	1	0	0	0	0	0.5	-	-	NS	NS	**								
27/6/72	19	2	3	2	10	7	6	10	6	9	4	7.0	***	6.4	NS	NS	NS								
26/9	2	1	1	2	4	3	1	3	1	1	2	1.9	*	1.8	NS	NS	NS								
16/11	2	1	0	1	2	2	1	1	1	1	1	1.2	***	0.98	*	NS	NS								

+ = Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Table 12

PROPORTION OF BROAD-LEAVED VOLUNTEERS IN THE PASTURE ON SIX OCCASIONS IN 1973-1974
(% dry weight)

	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)						Grand Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. on SR.	Hetero. Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15					17
18/6/73	13	14	37	21	12	10	10	21	25	22	28	19.3	**	**	NS
17/9	9	4	17	9	19	12	9	17	11	9	11	11.7	NS	NS	NS
26/11	34	3	24	9	38	30	27	31	25	8	7	21.7	**	**	NS
10/6/74	48	17	36	26	46	37	27	54	40	29	20	35.0	NS	NS	NS
2/9	31	10	25	18	26	27	22	25	27	13	18	22.2	NS	NS	NS
18/11	18	3	6	8	5	10	6	11	12	4	4	7.8	*	NS	NS

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Table 13

PROPORTION OF NARROW-LEAVED VOLUNTEERS IN THE PASTURE ON SIX OCCASIONS IN 1971-1972
(% Dry Weight)

Date	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)					Grand Hetero. mean			Sig. mean Reg. Coeff. on SR,	
	A	B	C	D	E	10	11	13	14	15	17	Adjt. treat. means	LSD 5% treat. means		Hetero. Reg. Coeff.
2/6/71	77	1	6	3	65	35	25	27	31	33	32	30.4	-	NS	*
8/9	46	0	6	2	54 [†]	21	19	21	17	24	24	21.0	-	NS	**
24/11	72	8	28	20	48	40	34	33	36	28	39	35.0	***	NS	NS
27/6/72	56	7	6	5	50	15	25	31	22	26	28	24.7	***	NS	NS
26/9	64 [*]	16	10	25 [*]	59	35	29	33	42	32	36	34.6	-	NS	**
16/11	77	23	23	34	44	44	34	37	47	36	41	40.1	***	NS	NS

[†] = Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Table 14

PROPORTION OF NARROW-LEAVED VOLUNTEERS IN THE PASTURE ON SIX OCCASIONS IN 1973-1974

(% dry weight)

Date	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)						Grand Hetero. mean	LSD 5% treat. means	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.			
	A	B	C	D	E	10	11	13	14	15					17		
18/6/73	42	18	3	7	7	43	24	25	21	17	24	25	22.5	***	11.4	NS	NS
17/9	8	19	7	7	7	7	8	12	10	6	11	11	9.6	*	7.9	NS	NS
26/11	57	65	45	54 ⁺⁺	32 ⁺⁺	32	43	45	46	44	66	60	50.6	*	18.5	*	NS
10/6/74	34	62	52	57	32	32	55	57	31	40	44	57	47.2	-	-	NS	*
2/9	38	66	45	48	37	37	53	66	43	33	48	38	46.6	NS	20.6	*	NS
18/11	63	65	53	46	32	32	53	61	57	45	44	48	51.7	-	-	*	**

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

⁺ = Significance of within-treatment regressions on stocking rate.

Although no data were collected on the botanical composition within either the broad- or narrow-leaved components it was obvious from visual observation that the proportion of barley grass, ripgut brome and capeweed was greatest at the lower stocking rates and *Vulpia* spp. and *Erodium* spp. at the higher stocking rates. During 1974, on no occasion was the botanical composition of either the broad-leaved component or the narrow-leaved component obviously different between pasture treatments at similar stocking rates.

8.4 Growth, crude utilization and availability of pasture

8.4.1 1971 Season

The significantly lower pasture production in treatment B and to a lesser extent in treatment D during the autumn (Figure 2) can be attributed to the poor growth rate of Wimmera ryegrass and Medea ryegrass respectively, as these components contributed over 90 per cent of total pasture production in their respective treatments (Table 5). However, total annual production was lowest in treatment D (Table 15), associated with this treatment having the smallest clover component during the spring period (Figure 1). Highest annual pasture production was recorded in treatments A and E (Table 15) associated with higher winter production from Yarloop subterranean clover in treatment A and higher spring production from Woogenellup and particularly Mt. Barker subterranean clovers[†] in treatment E (Figure 1).

The ranking of treatments based on the crude utilization of pasture was similar to that based on pasture growth rate in 1971 (Figure 2).

[†] The partitioning of the subterranean clover component into contributions by the Woogenellup and Mt. Barker subterranean clover varieties was only assessed by visual observation.




Figure 2. Mean rate of growth, mean rate of crude utilization and mean availability of pasture in all treatments during the experiment.

Table 15

ANNUAL PASTURE PRODUCTION FOR THE EXPERIMENTAL PERIOD
(kg ha⁻¹ year⁻¹ dry matter)

Date	Pasture treatment means				Stocking rate means (kg ha ⁻¹)						Grand Hetero. mean	Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. Coeff. on SR.		
	A	B	C	D	E	10	11	13	14	15					17	
1971	10885	7570	8850	7235	10325	9000	9035	8350	9350	9420	8685	8970	***	1236	NS	NS
1972	7230	6875	5575	5665	6240	5950	7010	6515	6155	5625	6640	6320	*	1178	NS	NS
1973	9300 ⁺	9190	8230	10090	8885	8215	9235	8400	9610	9435	9955	9140	-	-	*	*
1974	9305	9180	9250	9280	9555	9010	9350	9425	9355	9705	9055	9320	NS	1081	NS	NS
MEAN	9180	8200	7980	8070	8750	8040	8660	8170	8620	8550	8580	8440	**	597	NS	NS

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

+ = Significance of within-treatment regressions on stocking rate

The one major exception was in treatment A during the spring period when the crude utilization was significantly ($P < 0.01$) less than in the other four pasture treatments.

The availability of pasture on June 2 was significantly lower in treatment B ($P < 0.001$) and treatment D ($P < 0.01$) than in the other three pasture treatments and there was little difference in availability of pasture between these latter three pasture treatments (A, C and E). However, by September 8 (early spring), the availability of pasture in treatment A was significantly ($P < 0.001$) greater than in any other pasture treatment and this advantage continued for the rest of the growing season (Figure 2).

No within-treatment effect of stocking rate on pasture production was evident, although a significant ($P < 0.05$) mean (of all treatments) increase in pasture production with increased stocking rate was recorded for the period July 5 to September 8 (appendix Table 6). The mean (of all treatments) crude utilization of pasture significantly ($P < 0.01$ and $P < 0.05$) increased with increased stocking rate for the two periods June 2 to July 5 and July 5 to September 8 respectively, and also in all cumulative periods during the growing season (appendix Table 10).

Mean (of all treatments) pasture availability declined significantly ($P < 0.001$) with increased stocking rate on all sampling occasions in 1971 with some within-treatment regressions reaching significance ($P < 0.05$) on two occasions (appendix Table 14).

Only on one occasion (pasture growth during the period April 24 to July 5) was a significant ($P < 0.05$) interaction between pasture treatment and stocking rate recorded for growth, crude utilization or availability of pasture in 1971 (appendix Tables 6, 10 and 14).

8.4.2 1972 Season

Abnormal rains (170 mm) fell during the 1971/72 summer months (December 62, January 59 and February 49 mm, Table 2) resulting in a substantial germination of subterranean clover in all treatments and some growth of Victorian ryegrass and hybrid phalaris. No pasture measurements were taken but a considerable quantity of green pasture was available during this period. However, during March the clover plants died and all pastures dried off and new pasture growth was not evident again until the beginning of May.

Availability of pasture was extremely low during the first sampling period (May 1 to June 27) following almost complete utilization of pasture grown (Figure 2).

Most evident from Figure 2 is the substantially-higher growth rate of pasture in treatment A and, to a lesser extent, in treatment E during this period. This was associated with a dominant proportion of volunteer annual species in these two treatments compared to the dominance of ryegrass species in treatments B, C and D (Figure 1, Tables 5, 11 and 13). The different growth rate of pasture between treatments was not positively associated with the proportion of clover present in the pasture on June 27 as the largest mean clover component was recorded in treatment B rather than in treatments A or E (Table 9).

The extreme grazing pressure, particularly at the high stocking rates, resulted in very low levels of available pasture (Figure 2) during this period and significantly ($P < 0.05$) depressed pasture production was recorded with increased stocking rate in treatments C and D (appendix Table 7). However, over all treatments, no effect of stocking rate on pasture production or interaction between pasture treatment and stocking rate was recorded (appendix Table 7).

Lower pasture production in treatments C and D in the autumn-

winter period resulted in only marginally less total annual pasture production in these treatments (Table 15). However, higher utilization of pasture during the spring resulted in less available pasture in treatments C and D at the end of the growing season (Figure 2) although differences between treatments in pasture availability on November 16 did not reach statistical significance (appendix Table 14).

No influence of stocking rate on mean (of all treatments) pasture growth was evident in any period in 1972 but crude utilization in two periods and availability on all occasions were significantly affected by stocking rate (appendix Tables 7, 11 and 14).

8.4.3 1973 and 1974 Seasons

Rainfall for these two seasons was above the average of 638 mm (Table 2) and excellent pasture growth (particularly in comparison to 1972) occurred in all treatments (Table 15, Figure 2).

Few differences due to pasture treatment were recorded in pasture production during 1973 and 1974. Some association of lowered pasture production with a higher broad-leaved component in some treatments was evident but inconclusive. The uniformity of pasture production between treatments in 1974 was associated with a four-year trend (1971 to 1974) toward like botanical composition in all pasture treatments (Figure 1) with the exception of a persistent hybrid phalaris component in treatment E.

In 1973 a significant increase ($P < 0.05$) in pasture production was recorded with increased stocking rate for the spring and total annual pasture production in treatment A, and for the mean of all treatments in these two periods (appendix Table 8). No such relationship was obvious in 1974.

Pasture availability was marginally lower in treatment E during the autumn and winter of 1974 with a particularly marked depression

during the spring (Figure 2). This was associated with significantly ($P < 0.01$) higher crude utilization of pasture in treatment E during the spring.

Pasture availability decreased as stocking rate increased in all treatments on all occasions in 1973 and 1974 and regressions of mean (of all treatments) pasture availability on stocking rate were always significant ($P < 0.05$) with one exception on June 10, 1974 (appendix Table 15).

8.5 Subterranean clover seed production and reserves

Hard seed reserves of subterranean clover, measured in August 1973, were significantly higher in treatments A and B compared to the other three pasture treatments (Table 16). This was associated with high availabilities of clover herbage in treatment A throughout 1971 and 1972 and in treatment B throughout 1972 (Table 17). However, regressions of the 1973 hard seed reserves measured in each paddock on the proportion (% dry weight), production and availability of clover herbage for each sampling period/occasion in 1971 and 1972 were performed and the only significant variate was the availability of clover herbage measured on September 26, 1972.

In treatment D, there was a significant decrease in the 1973 hard seed reserves with increased stocking rate and this was the only occasion on which stocking rate could be significantly ($P < 0.05$) correlated with any subterranean clover seed parameter, within any pasture treatment or over all pasture treatments, during the experimental period.

The proportions of black seed (non-Yarloop) in the overall pool of hard seed reserves sampled in August 1973 in treatments A, B, C, D and E were 1, 4, 8, 5 and 21 per cent respectively with approximately equal proportions of black and white seed sampled above and below ground.

Table 17

AVAILABILITY OF SUBTERRANEAN CLOVER MEASURED ON SIX OCCASIONS IN 1971-1972
(kg ha⁻¹ dry matter)

Date	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)						Grand Hetero. Mean	Hetero. Adj. treat. means	LSD 5% treat. means on SR.	Sig. mean Reg. Coeff.		
	A	B	C	D	E	10	11	13	14	15					17	
2/6/71	180	5	10	5	65	30	60	65	70	50	40	55	***	48.6	NS	NS
8/9/71	730	25	30	5	170	240	195	290	220	120	85	190	***	177	NS	NS
24/11/71	995	460	410	30	1180	465	655	750	480	950	390	615	***	531	NS	NS
27/6/72	25	55	10	10	30	20	40	25	25	30	20	25	*	27.8	NS	NS
26/9/72	330	555	120	120	145	195	380	270	245	220	215	255	***	141	NS	NS
16/11/72	480	1780	930	695	1015	725	2005	1165	640	690	645	980	*	910	NS	NS

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Table 16

CLEAN SEED YIELD OF SUBTERRANEAN CLOVER SAMPLED ON FOUR OCCASIONS DURING
THE EXPERIMENT
(kg ha⁻¹)

Sampling Date	Pasture treatment means					LSD 5%
	A	B	C	D	E	
Hard seed August 1973	389	349	164	185***	139	94.6
Seed prod. below 1973	234	87	166	208	167	126.0
Seed prod. above 1973	57	31	46	54	43	25.2
Total seed prod. 1973	291	118	212	262	210	147.0
Hard seed August 1974	334	272	154	189	177	98.3
Seed prod. below 1974	-30	187	246	155	143	170.0
Seed prod. above 1974	85	55	74	41	66	32.5
Total seed prod. 1974	55	242	320	196	209	163.0

** = P < 0.01

+ = significance of within-treatment regression on stocking rate.

Table 17

AVAILABILITY OF SUBTERRANEAN CLOVER MEASURED ON SIX OCCASIONS IN 1971-1972
(kg ha⁻¹ dry matter)

Date	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)							Grand Hetero. Mean	Hetero. Adj. treat. means	LSD 5% treat. means on SR.	Sig. mean Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15	17					
2/6/71	180	5	10	5	65	30	60	65	70	50	40	55	***	48.6	NS	NS
8/9/71	730	25	30	5	170	240	195	290	220	120	85	190	***	177	NS	NS
24/11/71	995	460	410	30	1180	465	655	750	480	950	390	615	***	531	NS	NS
27/6/72	25	55	10	10	30	20	40	25	25	30	20	25	*	27.8	NS	NS
26/9/72	330	555	120	120	145	195	380	270	245	220	215	255	***	141	NS	NS
16/11/72	480	1780	930	695	1015	725	2005	1165	640	690	645	980	*	910	NS	NS

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Table 18

AVAILABILITY OF SUBTERRANEAN CLOVER MEASURED ON SIX OCCASIONS IN 1973-1974

Date	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)						Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. on SR.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17					
18/6/73	520	185	170 [†]	125 [*]	285 ^{**}	350	310	330	270	185	100	260	***	146	**	NS
17/9/73	1935	1285	985 ^{**†}	1115 ^{**†}	965	1805	1710	1130	1310	965	625	1260	**	543	***	NS
26/11/73	280	545	740	535	795	450	395	405	890	625	705	580	NS	486	NS	NS
10/6/74	285	110	100	130	115	65	50	105	205	235	220	150	*	124	**	NS
2/9/74	555	400	585	480	430	200	125	545	685	490	890	490	NS	452	**	NS
18/11/74	815	1060	1450	1390	1365	975	745	950	1410	1780	1440	1215	NS	608	*	NS

[†] = Significance of within-treatment regressions on stocking rate.

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

There was only a slight decline (c. 1%) in these proportions in the August 1974 sampling.

There was little between-year change in the hard seed reserves from 1973 to 1974, with the exception of a 22 per cent decline in treatment B (Table 16). The 'negative production' of below-ground seed in treatment A during 1974 would seem highly unlikely and must be attributed to sampling error, as there was no evidence of germination in the period August 1974 to January 1975 and the above-ground seed production was very similar to that in the other four pasture treatments (Table 16).

The data show that in both years in which the seed reserves were measured in January, by far the greatest proportion of seed present at that time was buried below ground[†] (Table 16) and appeared unrelated to treatment or stocking rate.

Regression analysis of clean clover seed produced in each paddock on seven pasture variates (measured in corresponding paddocks) was performed for the 1973 and 1974 data.

The pasture variates used were:

- (1) Availability of clover herbage at the end of winter.
- (2) Availability of clover herbage at the end of spring.
- (3) Mean availability of clover during the spring.
- (4) Proportion of clover (% dry weight) at the end of winter.
- (5) Proportion of clover (% dry weight) at the end of spring.
- (6) Mean proportion of clover (% dry weight) during the spring.
- (7) The production of clover herbage during the spring.

Clean seed production in both years was significantly correlated

[†] It is not to be implied that this was the actual ratio of seed produced above to that produced below during the growing season as no account has been taken of seed eaten or otherwise disappearing during the period August to January.

only with the availability of clover at the end of each respective spring according to the following equations

$$\text{1973 } y = 63.48 + 0.479x - 0.00024x^2, r^2 = 0.46, P < 0.001$$

$$\text{1974 } y = 24.98 + 0.188x, r^2 = 0.52, P < 0.001$$

where y = clean clover seed production (kg ha^{-1}) measured in 1973 and 1974 respectively

and x = availability of clover (kg ha^{-1}) measured on the last sampling occasion in 1973 and 1974 respectively.

It is interesting to note here that, although there appeared to be an increase in the amount of clover seed produced with increased availability of clover herbage at the end of the 1974 growing season, there was no recorded effect of stocking rate on clover seed production in 1974. This was despite the significant ($P < 0.05$) mean (of all treatments) increase in the availability of clover with increased stocking rate recorded on November 18, 1974 (Table 18).

8.6 Sheep liveweight data

The following points are evident from Figure 3.

8.6.1 General Trends

- a) The recorded pattern of liveweight change in the sheep is characteristic for the Mediterranean-type environment, with a decline in live weight during the summer and autumn (and in 1972, winter also) and increase during the winter and spring.
- b) Minimum, and maximum, live weights were recorded simultaneously in all treatments in all years.
- c) In the three years 1971, 1973, and 1974 minimum live weights were recorded during the same fifteen day, calendar period in each year (April 18 to May 3). In 1972, minimum live weights were not reached

until July 12 following the late seasonal break.

- d) The diminishing within-year magnitude of difference in live weight between pasture treatments over the experimental period (1971 to 1974).

8.6.2 1971

In comparison to the other four pasture treatments the mean live weight in treatment B declined more rapidly prior to the opening rains in April, and during May all sheep died from undernutrition at the highest stocking rate (17 ha^{-1}) in this treatment (appendix Table 29). Treatment B ranked significantly ($P < 0.001$) lowest in mean live weight on all weighing occasions in 1971 and on the first weighing in 1972 (January 19) prior to shearing. However, once new pasture growth occurred in April, liveweight gain was greater in treatment B than in treatment D.

Five distinct periods of differential liveweight change between pasture treatments were recorded:

- a) The initial period from the allocation of the sheep to their respective paddocks to the first paddock weighing (February 10). The decline in the liveweights in treatment B was the most significant aspect in this period.
- b) May 3 to June 23 - markedly superior liveweight gain was recorded in treatment A and, to a lesser extent, in treatments C and E.
- c) June 23 to August 24 - superior liveweight gain was only recorded in treatments A and E.
- d) August 24 to October 5 - substantial liveweight gain only occurred in treatment A.
- e) October 5 to November 11 - liveweight gain in treatment A was the lowest of all pasture treatments.

A significant effect of stocking rate on the mean (of all treatments) liveweight was first recorded on June 23 and an interaction between

stocking rate and pasture treatment was recorded on November 11 (heterogeneity of regression coefficients was significant ($P < 0.001$)). This latter interaction was manifested in the different pattern of change (with time) in the value of the regression coefficient of the higher ranked pasture treatments (A, E and to a lesser extent C) compared to the lower ranked treatments (B and D) after August 24. The higher ranked pasture treatments recorded a marked maximum depression of live weight with increased stocking rate on August 24 compared to a less marked maximum on October 5 in treatments B and D.

8.6.3 1972 and 1973

Significant aspects evident in the mean liveweight response during 1972 and 1973 were:

- a) The more rapid liveweight gain in treatments A and B compared to the other pasture treatments following the late start to the growing season (May 1, 1972). This superior liveweight gain in these two pasture treatments continued until the end of September. The mean live weight in treatment A was also significantly ($P < 0.05$) higher than in the other four pasture treatments on August 16 and September 28, 1972.
- b) Treatment B increased in rank (liveweight basis) during the 1972/73 season from the lowest ranking on March 8, 1972 to the highest on January 17, 1973.
- c) Following the seasonal break in 1972, treatment D ranked the lowest and treatment C the second lowest, in mean live weight throughout 1972 and 1973. The mean live weight in treatment D was significantly ($P < 0.05$) lower than the mean live weight in all other pasture treatments on two occasions (November 29, 1972 and January 17, 1973) and significantly ($P < 0.05$) lower than the higher ranked two pasture

treatments (A and B) only on some other occasions.

- d) A significant ($P < 0.05$ but usually $P < 0.001$) effect of stocking rate on the mean (of all treatments) live weight was recorded on all occasions in 1972 and 1973 except on November 20, 1973. In all cases increased stocking rate resulted in lower mean liveweights of the sheep.
- e) Although large differences existed between pasture treatments in the magnitude of the regression coefficient on many occasions in 1972 and particularly in 1973 (i.e. January 17 to July 17, 1973), significant differences ($P < 0.05$) were only recorded on three occasions i.e. April 18, 1972, August 21, 1973 and October 3, 1973, and in all cases, the decline in live weight with increased stocking rate was greatest in treatment D.

Also in treatment D, the maximum value of the regression coefficient was not reached until January 17, 1973 after the 1972 season although maxima were attained on September 28, 1972 in the other four pasture treatments. However in 1973, the pattern of change in value of the regression coefficient with time was similar in all pasture treatments.

8.6.4 1974

No significant ($P > 0.05$) difference in mean live weight between pasture treatments was recorded in 1974. Significant effects of stocking rate on live weight were only recorded (and then only in some pasture treatments) on three occasions i.e. immediately before, at, and immediately after the occurrence of minimum live weight (May 1, 1974). The value of the regression coefficient was again highest in treatment D for most of the season although a significant ($P < 0.05$) interaction between stocking rate and pasture treatment on live weight was only recorded on one occasion (June 6, 1974).





Figure 3: Mean live weight and linear regression coefficient of live weight on stocking rate for each pasture treatment.

8.7 Annual wool production

Significant interactions between pasture treatment and stocking rate on the clean wool produced per head and per hectare were recorded in both 1971 and 1973 but not in 1972 and 1974 (Table 19, Figures 4 and 5). With the exception of the mean (of all treatments) clean wool produced per head in 1974, stocking rate significantly ($P < 0.001$) affected the mean (of all treatments) clean wool produced, both on a per head and per hectare basis, in all years. The slopes of the mean regression lines were negative in the case of annual clean wool produced per head and positive in the case of annual clean wool produced per hectare (Figures 4 and 5).

The significance of individual within-treatment regressions of annual clean wool produced on stocking rate are given in Table 19. In no case could a polynomial relationship of higher order than a linear relationship be statistically justified over the range of stocking rates used in this experiment.

The mean annual clean wool produced was considerably lower in treatments B and D compared to the other treatments in 1971 (Table 19). The statistical significance of the difference in treatment means cannot be calculated due to the interaction between pasture treatment and stocking rate in 1971.

In 1972 treatment D produced the least, and treatment A the most, clean wool. The difference between these two treatments and the next ranked pasture treatment (above and below respectively) was significant ($P < 0.05$) in both cases.

Clean annual wool production was again lowest in treatment D in 1973, but there was no difference in production recorded between pasture treatments in 1974 (Table 19).

Table 19

ANNUAL CLEAN WOOL PRODUCTION
(kg)

	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)					Grand Mean	Hetero. LSD Adj. treat. means	5% mean Reg. Coeff. on SR.	Sig.	Hetero. Reg.			
	A	B	C	D	E	10	11	13	14						15	17	
1971	per head	3.6	2.4†	3.7	2.9	3.8	3.6	3.7	3.4	3.3	3.1	3.1	3.1	3.3	-	***	*
	per ha	* 47	**† 31†	* 49	37	*** 50	36	40	44	46	47	53	43	-	-	***	***
1972	per head	* 3.7	** 3.3	*** 3.3	* 3.0	* 3.4	3.8	3.9	3.4	3.2	3.1	2.7	3.3	***	0.26	***	NS
	per ha	48	43	*** 43	38	45	38	42	45	45	47	52	44	***	4.0	***	NS
1973	per head	5.7	5.7	* 5.6	5.2	5.6	5.9	5.7	5.6	5.5	5.4	5.3	5.6	-	-	***	*
	per ha	** 76	** 75	* 73	** 69	** 74	59	63	72	76	81	89	74	-	-	***	*
1974	per head	4.5	4.5	4.6	4.6	4.6	4.8	4.6	4.7	4.5	4.5	4.5	4.6	NS	0.29	NS	NS
	per ha	*** 60	* 60	** 61	** 61	** 62	48	51	60	62	67	77	61	NS	4.0	***	NS

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

† = mean of 5 lowest stocking rates only.

+ = significance of within-treatment regressions on stocking rate.




Figure 4. Relationship between clean wool produced per head and stocking rate.
 NA=not available NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001




Figure 5. Relationship between clean wool produced per hectare and stocking rate.
 NA=not available NS: P>0.05 * P<0.05 ** P<0.01 *** P<0.001

The mean (of all treatments) quantity of clean wool produced was very similar in 1971 and 1972 and maximum annual production occurred in 1973 (Table 19).

8.8 Fleece characteristics

No significant interaction between pasture treatment and stocking rate was recorded for any of the fleece characteristics in any year (Tables 20 and 21). Significant ($P < 0.001$) differences between pasture treatment means were recorded for staple length and fibre diameter and ($P < 0.05$) for yield in 1971 only and not in any of the following three years (Tables 20 and 21). Treatments ranked (on a staple length and fibre diameter basis) as follows, with differing subscripts indicating a significant ($P < 0.05$) difference between pasture treatment means.

Staple length - $E_a > C_a > A_{ab} > D_b > B_c$

Fibre diameter - $A_a > E_{ab} > C_b > D_c > B_c$

Highly significant ($P < 0.001$) mean (of all treatments) regressions (negative in slope) of staple length and fibre diameter on stocking rate were recorded in 1971 and 1972 and with less significant correlation in 1973. Significant negative correlations between stocking rate and both yield in 1972 and crimp frequency in 1973 were also recorded (Tables 20 and 21).

Fibre diameter increased and both crimp frequency and yield decreased between years from 1971 to 1974 (Tables 20 and 21). In each case, the mean (of all treatments) regression of fleece characteristic on year was significant ($P < 0.05$).

8.9 Seasonal wool production

Season had a far greater influence on wool growth rate than did

pasture treatment. This is clearly demonstrated in Figure 6 by the amplitude of fluctuations in the changing rate of wool growth compared to the magnitude of between-treatment differences in wool growth rate during the experimental period.

During 1971 and 1972 the mean wool growth rate varied in a similar pattern to the mean liveweight change in the sheep (Figures 3 and 6) with periods of maximum rate of wool growth approximately coinciding with periods of maximum liveweight gain in the sheep.

In 1971 the mean live weights of the sheep increased throughout the spring and early summer but the mean wool growth rate declined from the end of October. However in 1973 and 1974 the mean wool growth rate declined from the beginning of August and the beginning of July respectively with a plateau and peak in mid-November and mid-December respectively. In both these years (1973 and 1974) this decline in wool growth rate during the spring occurred despite constant liveweight gain of the sheep during this period (Figure 3).

With large, seemingly random, fluctuations in the value of the regression coefficient (wool growth rate change per unit increase in stocking rate) with time (Figure 6), significant ($P < 0.05$) Stocking rate x Pasture treatment interaction in respect of wool growth rate was only recorded in the first three growth periods in 1971 (Figure 6). In these three periods, wool growth rate in treatments E and A was less depressed by the higher stocking rates than in treatments C, B and D (the latter three treatments being ranked on increasing absolute value of the regression coefficient). In all other growth periods in 1971 and 1972, significant ($P < 0.01$) and usually $P < 0.001$) differences in the mean wool growth rate between pasture treatments were recorded (Figure 6). The only two exceptional periods were during the mid-summer of 1971/72 and 1972/73.

However in 1973 and 1974, only in two consecutive growth periods, was a significant difference in wool growth rate between pasture treatments recorded (during the spring of 1973).

The ranking of pasture treatments based on mean wool growth rate was consistent with the ranking of treatments based on mean live weight on most occasions in 1971 and 1972. This applied in the case of treatments D and B even though, on the basis of liveweight gain, the order was reversed. Treatments E and B during most of 1971 and treatments A and D in 1972, ranked highest and lowest respectively in terms of both wool growth rate and live weight (Figures 3 and 6).

In all growth periods in 1971 and 1972 the mean (of all treatments) depression of wool growth rate with increased stocking rate was significant ($P < 0.05$ and usually $P < 0.001$) but in 1973 and 1974 a similar relationship was only recorded significantly ($P < 0.05$) in some growth periods during the late winter and spring (Figure 6).

Table 20

FLEECE CHARACTERISTICS 1971 AND 1972

	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)					Grand Mean	Hetero. Adj't. treat. means	LSD 5% treat. means SR,	Sig. Mean Reg. Coeff.	Hetero. Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15						17
<u>1971</u>																
Staple leng. (cm)	9.5	†8.3	9.8	9.1	9.9	9.5	9.8	9.5	9.1	9.3	9.2	9.3	***	0.44	***	NS
Fibre diam (µm)	20.4	†18.2	19.7	18.7	20.2	20.2	19.8	19.9	19.5	19.0	18.8	19.4	***	0.69	***	NS
Crimp per cm	3.4	†3.7	3.7	4.1	3.5	3.7	3.5	3.6	3.6	4.0	3.7	3.7	NS	0.49	NS	NS
Yield (%)	75.0	†76.4	79.0	77.1	77.6	77.3	77.0	78.1	76.3	75.7	77.5	77.0	*	2.46	NS	NS
<u>1972</u>																
Staple leng. (cm)	8.5	8.4	8.6	8.3	8.7	8.7	9.3	8.8	8.3	8.2	7.8	8.5	NS	0.63	***	NS
Fibre diam (µm)	20.6	20.3	20.0	20.0	20.2	21.0	21.0	20.6	19.9	20.0	18.7	20.2	NS	0.92	***	NS
Crimp per cm	3.5	3.1	3.3	3.4	3.3	3.5	3.2	3.2	3.3	3.3	3.5	3.3	NS	0.34	NS	NS
Yield (%)	73.2	73.8	76.1	75.6	75.0	76.1	75.7	75.5	73.5	72.8	74.4	74.7	NS	2.25	*	NS

NS = P>0.05 * = P<0.05 ** = P<0.01 *** = P<0.001

† = Significance of within-treatment regressions on stocking rate

‡ = Mean of 5 lowest stocking rates only.

Table 21

FLEECE CHARACTERISTICS 1973 AND 1974

	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)							Grand Mean	Hetero, LSD Adj't, treat, means	Sig. mean Reg. Coeff. on SR.	Hetero, Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15	17					
<u>1973</u>																
Staple leng. (cm)	10.0	10.2	9.8	9.8	10.1	10.2	10.3	10.2	9.7	9.8	9.6	10.0	NS	0.55	*	NS
Fibre diam (µm)	23.0	23.4	22.6	22.7	22.9 [†]	23.8	23.1	23.1	22.6	22.8	21.9	22.9	NS	0.95	**	NS
Crimp per cm	3.2	3.2	3.1	3.0	3.1	3.3	3.1	3.1	3.2	3.1	3.0	3.1	NS	0.20	**	NS
Yield (%)	73.3	75.0	75.4	74.4	74.3	75.7	74.4	75.5	73.9	73.6	74.3	74.5	NS	2.78	NS	NS
<u>1974</u>																
Staple leng. (cm)	9.8	9.9	9.8	9.8	9.6	9.8	9.8	10.2	9.6	9.5	9.8	9.8	NS	0.66	NS	NS
Fibre diam (µm)	23.3 ^{**}	24.2	23.2	24.0	23.9	24.6	23.7	24.0	22.9	24.0	23.2	23.7	NS	1.10	NS	NS
Crimp per cm	3.1	2.9	3.0	3.1 [*]	2.9	3.1	3.0	3.1	3.1	2.9	2.9	3.0	NS	0.23	NS	NS
Yield (%)	71.1	73.1	73.2	73.5	73.2	73.5	72.3	74.0	72.7	72.1	72.5	72.8	NS	2.72	NS	NS

NS = P>0.05 * = P<0.05 ** = P<0.01 *** = P<0.001

† = Significance of within-treatment regressions on stocking rate





Figure 6: Mean clean wool weight and linear regression coefficient of clean wool weight on stocking rate for each pasture treatment.

DISCUSSION AND CONCLUSION

9. DISCUSSION AND CONCLUSION

9.1 Animal production in relation to botanical composition

It is necessary to first refer to the fact that treatment A did not have the tillage and seedbed preparation, or the sowing of new species. While it is acknowledged that in some respects this has a confounding influence, the pasture treatments are best regarded as 'mini systems' which can be validly extrapolated to the farm situation. In this respect, the various pasture treatments or pasture systems, with superimposed stocking rate treatments, are quite valid.

9.1.1 The importance of subterranean clover

The outstanding result from this experiment is that the renovation of the existing volunteer annual grass - Yarloop subterranean clover pasture by the tillage, seedbed preparation and sowing of the four grass - subterranean clover mixtures did not result in either increased live weight or wool production from the grazing wethers during the set-stocked experimental period of four years. In fact, lower pasture and animal production occurred in the three ryegrass treatments during the first and, to a lesser extent, the second and third years of set stocking.

The differences in productivity can be attributed to the extensive differences in availability of clover that existed between pasture treatments in 1971 and, to a lesser extent, in 1972 and 1973, following the renovation and pasture management procedures carried out in 1970.

It is difficult to ascribe animal productivity to the quantity and quality of particular botanical components of grazed pasture and the quantitative and qualitative intake of pasture by the grazing animal may be related (Conrad 1966). The most important single determinant of

ruminant production at a particular time is the quantity of feed being harvested by the animal, and this is very often the immediate limiting factor to animal production. Demonstration of qualitative differences can only be accomplished when allowance has been made for differences in quantity of feed eaten (Lambourne 1969).

In the comparison of treatments cited below, quality and quantity of pasture are confounded because total pasture production was usually higher in those treatments with a higher clover component.

Nevertheless, the data that have been presented, strongly implicate the availability of the subterranean clover component as being the dominant factor in the determination of animal production in this experiment. It may be argued that the differences in animal performance in any of the treatment comparisons cited below, may be due to the higher quality of the non-legume component. However this is unlikely to be true as there is little evidence in the literature for substantially-superior nutritional value of any of the volunteer or sown grass species comprising the pastures in this experiment. For example, McIvor and Smith (1973c) reported little difference in the dry matter content, nitrogen content of leaves and *in vitro* digestibility of a number of temperate annual pasture species including ripgut brome, barley grass and annual ryegrass. Thus during the autumn of 1971, for example, it is unlikely that the superior liveweight gain recorded in treatment A was due to the higher nutritional value of the barley grass-dominant pasture compared to the pasture comprising dominant ripgut brome and hybrid phalaris in treatment E or Victorian ryegrass in treatment C, but rather to the higher proportion and availability of subterranean clover. Certainly there was no substantial difference between these pasture treatments in pasture production, crude utilization or availability of pasture during this period. This example quoted is the only one in which superior

quality *per se* of the clover herbage was clearly demonstrated because equivalent levels of production, utilization and availability of total pasture were recorded.

Further examples implicating the availability of clover as the main determinant of animal production in this experiment are as follows:

- (1) The proportion and availability of clover, together with the total availability of pasture, were significantly greater in treatment A during the winter of 1971, and only in this treatment did the mean live weight increase substantially during this period.
- (2) Comparing the three ryegrass treatments B, C and D during the growing season of 1971, the lowest availability of clover was recorded in treatment D and this was associated with the lowest liveweight gain being recorded in this treatment.
- (3) During the winter and spring of 1972 the rapid liveweight gain in treatment B was associated with maximum proportion and availability of clover being recorded in this treatment. The increased performance of the sheep in treatment B during 1972 is demonstrated by the change in rank (liveweight basis) of this treatment, from the lowest on March 8, 1972 to the highest on January 17, 1973 and this was associated with this treatment having the greatest between-year increase in the size of the subterranean clover component of all the pasture treatments from 1971 to 1972.
- (4) The lower liveweight gain during the winter of 1972 in treatments C, D and E was associated with lower availabilities of subterranean clover during this period.
- (5) The decline in live weight with increased stocking rate was greatest in treatment D throughout 1973 and 1974 and only in this treatment was a decrease in the proportion of clover with increased stocking rate recorded.

9.1.2 The role of other species

Apart from the role of subterranean clover, no other consistent relationship between the size or presence of a particular pasture component and animal performance was evident in this experiment although two isolated instances are worthy of note.

- (1) The greater liveweight loss in treatment B, prior to the opening of the season in 1971, can only be explained by the poor nutritional value of the available feed in this treatment over the summer period. The availability of pasture at the end of the 1970 growing season was similar in treatments B and E and even lower in treatment A.

Reasons for this poor quality of feed in treatment B are possibly:

- a) A lower clover component in the dry summer residue resulting from suppression of this component by the Wimmera ryegrass during the spring of 1970, even though an attempt had been made to prevent this happening by the strategic grazing of the pasture at that time.
- b) Lower nutritional value of the Wimmera ryegrass itself, due to a lower number of seed heads being available to the sheep in comparison to the previously ungrazed Victorian and Medea ryegrasses. However, these latter two treatments also had higher availabilities of pasture at the end of the 1970 growing season.

In comparison to treatments A and E the Wimmera ryegrass may have been of lower nutritive value than the dominant barley grass and ripgut brome components that comprised treatments A and E respectively. Radcliffe and Cochrane (1970) reported that in Wimmera ryegrass the digestibility declined at the rate of 0.9 units per day in the month following flowering compared to an average of 0.3 to 0.5 units per day for phalaris,

brome grass, silver grass, barley grass, Yarloop subterranean clover and barrel medic. They also recorded a low crude protein content in Wimmera ryegrass in mid-December.

- (2) During 1971 and 1972, the decline in rate of liveweight gain in treatment A during mid to late spring and marked decline in wool growth rate during the spring of 1971 compared to the other pasture treatments could be attributed to an excessive production of, and aggravation by, barley grass inflorescences (Campbell *et al.* 1973).

9.2 Persistence of sown species

9.2.1 Grasses

The almost total failure of Medea ryegrass to demonstrate any perennial characteristics under continuous grazing in this experiment was both unexpected and difficult to explain and repudiates the original claims of increased individual plant persistence by this cultivar, at least under the conditions imposed in this experiment. Certainly, Medea ryegrass demonstrated adequate seed production and an ability (marginally greater than Wimmera and Victorian ryegrasses) to regenerate from self-sown seed following two summers in this environment.

However, as has frequently been demonstrated in the past, the behaviour of plant species under continuous grazing may differ from that demonstrated under cutting or short, periodic grazing regimes (Davidson 1969).

Subsequent to the completion of this experiment, reports of wide variability in the summer dormancy and persistence of Medea ryegrass have been received in South Australia (Higgs, personal communication).

An experiment (Gibson, unpublished) carried out in 1971 and 1972 on Kangaroo Island to define the best first year grazing management

practice for Medea ryegrass demonstrated that this species could exhibit a marked degree of perenniality when intermittently grazed during the spring and summer in this environment, but not when under continuous grazing by sheep.

The lack of persistence of Victorian ryegrass over more than three summers in this environment agrees closely with the findings of Carter and Day (1970), on Kangaroo Island. These authors attributed the decline in the perennial ryegrass component, in their experiment, to the dry spring and summer conditions. The death of the Victorian ryegrass plants in this experiment occurred predominantly over the 1972/73 summer which also followed the driest spring and summer encountered during the experimental period: the total rainfall during the 1970/71, 71/72, 72/73 and 73/74 spring/summer periods being 172, 354, 91 and 247 mm respectively.

The influence of increased stocking rate in hastening the between-year decline in density of Victorian ryegrass plants, and the proportion of this species in the pasture, was clearly demonstrated in this experiment. Decreased persistence of perennial ryegrass with increased stocking rate has also been reported by Cameron and Cannon (1970) in Victoria.

In this Kangaroo Island experiment the species failed to persist even at the lowest stocking rate and it seems likely that the dry spring/summer conditions, possibly in combination with continuous grazing during this period, probably caused the lack of persistence of this species in this environment.

The high degree of persistence of the hybrid phalaris during the experimental period, irrespective of stocking rate, is in marked contrast to that of the ryegrass species and demonstrates the ability of this species to persist under variable management and environmental influences. This finding is in accordance with that of Hutchinson (1970) at Armidale,

New South Wales, in a different environment and with that of Reed (1974) in a more similar environment at Glenormiston, Victoria. Both of these workers also demonstrated an interaction between survival of perennial grass species and stocking rate in their experiments.

Although stocking rate did not influence the density of hybrid phalaris plants, it did influence their size and also the proportion of this species in the pasture, however, statistical significance could not be demonstrated during the experimental period. Visual appraisal of the pastures during the spring of 1975 and 1976 (after intermittent grazing during these two years) revealed a continuation of the between-year trend towards increasing plant size and proportion of the hybrid phalaris in the pasture. In some paddocks the proportion was estimated to be between 30 and 40 per cent of the total available pasture. This highlights the problem of evaluating pasture species only over a relatively short period (4 years) when the long term (at least 10 years) changes in botanical composition may be important.

A further grazing experiment (Gibson, unpublished) in which the same hybrid phalaris has been compared with perennial ryegrass and Demeter fescue under grazing at three stocking rates in a higher rainfall environment, has since demonstrated high pasture production from the hybrid phalaris at the higher plant densities obtained in that experiment. However there was no advantage over the other pasture species in terms of animal production.

9.2.2 Subterranean clover

The partial replacement only of Yarloop with Mt. Barker and Woogenellup subterranean clovers in the renovated treatments in 1970, proved unsuccessful in maintaining a low proportion of Yarloop subterranean

clover in the overall clover component after a period of four to five years.

Although, in 1970, the growth and seed set of the introduced subterranean clovers was suppressed to some degree by the vigorous growth of the ryegrass in treatments B, C and D, this did not occur in treatment E, yet Yarloop successfully invaded this treatment. This is highlighted by the level of hard seed reserves present in August 1973 in treatment E (c. 140 kg ha⁻¹) of which only 21 per cent was non-Yarloop seed. However, as the level of subterranean clover seed reserves was not measured in 1970, it is uncertain to what extent any initial reserves were important in enabling the rapid invasion of Yarloop into the pastures to take place in subsequent years.

Beale (1974) found that germination of hard and soft seed of Yarloop from an initial reserve on Kangaroo Island, continued at a very high level in the second and third years and only began to fall off in the fourth year. This is in contrast to the findings of Donald (1959) and Taylor (1972) who found that the majority of subterranean clover seed germinated in the first year after the seed was set and that the level of regeneration fell rapidly in subsequent years. However, these latter two workers did not include Yarloop in their experiments and Beale (1974) suggests that Yarloop has a greater capacity for continual regeneration from seed than has been observed in Geraldton, Dwalganup, Bacchus Marsh, Mt. Barker and Tallarook subterranean clovers.

In this report experiment the level of hard seed reserves were similar in treatments A and B by August 1973 indicating that any decline

treatment B, following the pasture renovation performed in this treatment, were only temporary.

possibility of replacing Yarloop with
nean clover are presently in progress on

Kangaroo Island. Particular attention is being paid to minimizing the germinable pool of Yarloop seed, prior to the sowing of new cultivars. Various tillage practices, such as mouldboard ploughing to deep-bury the seed and/or repeated cultivation to kill successive germinations of Yarloop are being evaluated (Beale *et al.* 1976).

The lack of evidence for the influence of stocking rate on the hard seed reserves and the seed production measured in 1973 and 1974 in this experiment, indicates that adequate levels of Yarloop seed reserves can be maintained, irrespective of the level of set stocking; at least over the time period and range of stocking rates used in this experiment.

The proportion of clover in the pasture on many occasions and particularly on the final sampling occasion in the spring of each year (except 1972), and the availability of pasture on nearly all sampling occasions during the experiment, were respectively positively and negatively correlated with stocking rate. Hence the net influence of stocking rate on the availability of clover herbage is dependent on the degree to which stocking rate influences both the proportion of clover in the sward and the total availability of pasture. In 1971 and 1972, there was no evidence of stocking rate influencing the availability of clover on any occasion and in 1973 and 1974, negative and positive relationships were recorded respectively.

In 1973 and 1974, only c. 50% of the variability in clover seed production was accounted for by the availability of clover at the end of the growing season. This is not surprising considering the possible variability in both the seed production and clover availability data due to sampling. This latter point is highlighted by the anomolous result obtained in 1974 when clover seed production was not influenced by stocking rate even though the former was related to the availability of clover, and stocking rate influenced the availability of clover.

Nevertheless, it is clear from both the pasture production data and the data on clover seed production and reserves that Yarloop is extremely well adapted to this environment with its outstanding growth during the winter and early spring and with its ability to set copious supplies of seed when grazed at both high and low stocking rates.

9.3 Invasion of non-leguminous species

The invasion of the renovated treatments B, C and D by volunteer annual species proceeded concurrently with the decline in the ryegrass component, and by 1974 obvious changes in botanical composition due to stocking rate were evident. The higher proportion of ripgut brome, capeweed and barley grass at the lower stocking rates, and *Vulpia* and *Erodium* spp. at the higher stocking rates, agrees with the findings of other workers (Sharkey *et al.* 1964, Davies 1966, Carter 1968b).

The occurrence of cluster clover and winter grass at the highest stocking rates as reported by Carter (1968b) and Brown (1976a) was not evident. However, it is possible that the effective grazing pressure (in addition to the actual stocking rate) at the highest stocking rate in this experiment was lower than in the above two experiments where there was need for regular supplementary feed in Brown's experiment and sheep deaths at the highest stocking rate in Carter's experiment. In this report experiment, sheep that died from undernutrition were from mainly one treatment (B) and these deaths occurred in only one year (1971).

The dominance of ripgut brome in the absence of grazing in treatment E during 1970 supports the observations of the Californian workers (Talbot and Biswell 1942, Jones and Evans 1960) and also those of Rossiter and Pack (1956) in Western Australia.

9.4 Interaction of stocking rate and botanical composition in the determination of animal production

It has been clearly demonstrated that the differences in animal production recorded in this experiment involving various pasture treatments and stocking rates were usually associated with the qualitative and quantitative presence of subterranean clover in the pasture. The presence or absence of other species (whether sown temperate, perennial or annual species, or volunteer, broad-leaved or narrow-leaved annual species) could not consistently be associated with changes in animal productivity in this experiment. The marked invasion of Yarloop into the renovated pasture treatments occurred at all stocking rates and, although the availability of clover may be influenced by stocking rate within any one season, in the longer term the presence of Yarloop appears independent of stocking rate. Hence the excellent adaptation of Yarloop in this environment, with its rapid growth during the winter and early spring and its ability to set copious supplies of seed with potential for germination over a number of years, seems to have guaranteed these pastures a degree of quantitative and qualitative independence of stocking rate. It therefore is important to state, that in the long term, there appears to be no obvious interaction of stocking rate and botanical composition in the determination of animal production in this environment. This is not to say that no such interaction occurred in this experiment (as it obviously did in 1971) but the difference in botanical composition was primarily a result of the pasture renovation practice rather than as a direct result of imposed stocking rates. The obvious effects of the renovation practice in terms of botanical composition were only evident in the short term.

The short term interaction of pasture treatment and stocking rate in determining animal production is best seen by comparing the quantity

of clean wool produced per hectare in 1971 in treatments A and D. The absence of a significant component of Yarloop in treatment D during 1971 and most of 1972 resulted in an almost constant quantity of clean wool per hectare being produced over the range of stocking rates. Only in 1973 was there a significant increase in clean wool produced per hectare with increased stocking rate. In contrast, maximum clean wool per hectare was produced at the highest stocking rate in treatment A in all years with no indication (except perhaps in the poorest pasture year of 1972) that the potential maximum clean wool per hectare had been attained. However, in 1974 there was no difference between these pasture treatments in the relationship of clean wool produced per hectare and stocking rate; a result of a significant legume component comprising the pastures of both pasture treatments.

9.5 Wool production and fleece characteristics

Clear evidence has been produced for an inverse relationship between the clean fleece weight and stocking rate of the wethers in this experiment. This result is to be expected in the light of previously reported experiments with wethers (Arnold and McManus 1960, Cannon 1969, Carter and Day 1970, Brown 1976) and with ewes (Davies 1962, Morley *et al.* 1969). The reduction in fibre diameter and staple length with increased stocking rate, as recorded in this experiment, has also been reported in previous experiments (Sharkey *et al.* 1962, Cannon 1972, Langlands and Bennett 1973, Lipson and Bacon-Hall 1974, White and McConchie 1976, Brown 1976b). However, McManus *et al.* (1964) in the Canberra environment, found no significant effect of stocking rate on the fibre diameters, staple lengths and subjective characteristics of Merino fleeces. However these workers did observe stocking rate effects on the seasonal changes in fibre diameter and staple length during their

experiment. Similar seasonal trends have been recorded by Roe *et al.* (1959) and Sharkey *et al.* (1962).

The observed decrease in staple crimp frequency with increased stocking rate in 1973, and the lack of response of this characteristic to stocking rate in other years in this experiment, is in contrast to the positive relationship reported by White and McConchie (1976), Cannon (1972) and Langlands and Bennett (1973).

Cannon (1972) reported both increases and decreases in yield of clean wool with increased stocking rate (depending on the level of superphosphate application in his experiment) while Langlands and Bennett (1973) and White and McConchie (1976) reported no relationship. However in this report experiment, a negative relationship was recorded in 1972 which is in accordance with the findings of Brown (1976b) who recorded a similar relationship in four consecutive years.

Although no significant interaction between pasture treatment and stocking rate was recorded for any of the fleece characteristics in any year, significant differences between pasture treatments were recorded in this experiment in 1971.

There appears to be no experiment reporting a significant effect of pasture type on the fleece characteristics of grazing sheep. In one study reported from Canberra, A.C.T. (McManus *et al.* 1964) in which the fleece characteristics of Merino wethers grazing either a phalaris-subterranean clover pasture or an annual pasture comprising subterranean clover, *Bromus* spp. and barley grass were compared, no effect of pasture type on any fleece characteristic was recorded. However, large differences in the proportion and availability of the pasture legume component did not exist in that experiment and hence the result is not in discord with this Kangaroo Island experiment.

In most experiments reported, the fleece characteristics of the

grazing sheep have been insensitive, not only to botanical composition but also to pasture management practices (other than stocking rate). For example, isolated instances only (and then in only one year) have been reported where the level of superphosphate application (Cannon 1972, White and McConchie 1976) or grazing management (Brown 1976b) have significantly affected any fleece characteristic.

9.6 Efficiency of high stocking rates

There is no doubt that the use of higher stocking rates on the annual-type pasture typical of the Mediterranean-type climate in southern Australia will result in increased animal production, particularly wool production as recorded in this experiment (Lloyd Davies and Humphries 1965, Carter 1965, Carter and Day 1970, Carter 1973, Brown 1976b).

The efficiency of wool production in relation to pasture production and rainfall, was clearly dependent on stocking rate (Table 22) and this experiment highlights the biological and economic inefficiency of low stocking rates. The indices of total clean wool production, in relation to rainfall, show a close parallel with that for pasture production, and emphasize the improved efficiency of water use at higher stocking rates. The percentage crude utilization of green pasture ranged from 57% to 73% at the lowest and highest stocking rates respectively (Table 22). Total annual crude utilization of pasture (including the pasture utilized or otherwise disappearing over the summer months) would be much higher. Carter (1973) reported total annual crude utilization values of 96.2% at a stocking rate of 22.2 sheep ha⁻¹ in his stocking rate experiment on annual pasture at the Waite Institute, South Australia. Those stocking rates in which a crude utilization value of 82% was recorded were perfectly safe resulting in no undue stress on the experimental wethers under the

conditions of his experiment.

This experiment has demonstrated an approximate linear response of wool production to increase in stocking rate up to the maximum of 17 wethers ha⁻¹ imposed in this experiment without any obvious adverse effects on the productivity of the grazed pastures. This is not to say that all pastures in this environment should be stocked at this level as allowance must be made for the additional requirements of breeding sheep and young growing sheep. In addition, the social and risk factors associated with high stocking rates must also be taken into account (Carter and Day 1970, McArthur 1970, McArthur and Dillon 1971) and it must be realized that probably three out of the four years of set stocking in this experiment, experienced above-average seasonal rainfall.

9.7 Conclusion

Although one cannot extrapolate the results of this relatively short-term experiment to the longer-term without some degree of uncertainty, this experiment has indicated that in the past undue emphasis may have been placed on minimizing the contribution of many of our volunteer annual grasses and herbs and attempting to stabilize botanical composition by sowing temperate perennial species rather than ensuring that an adequate legume component is present.

Provided that the problem of clover-induced infertility in ewes can be overcome, either by sheep selection and/or managerial means or by the replacement of Yarloop subterranean clover with an equally-productive and well adapted legume, high animal production should be maintained from the annual pastures in this environment.

Certainly, change in botanical composition resulting from various stocking rates *per se* is no cause for alarm, but rather it indicates the

weaknesses of our sown species and the strengths of many of our naturally-occurring volunteer annual species.

Table 22

MEAN PRODUCTION DATA AND INDICES OF CLEAN WOOL PRODUCED IN RELATION
TO PASTURE PRODUCTION AND APRIL-NOVEMBER
RAINFALL, PARNDANA EAST 1971-74

Stocking Rate (sheep ha ⁻¹)	Pasture grown kg ha ⁻¹ D.M.	Annual green C.U. kg ha ⁻¹ D.M.	Proportion green pasture utilized during growing season %	Clean Wool grown kg ha ⁻¹	Wool grown kg/1000 kg DM.	Wool grown kg ha ⁻¹ / 100 mm Ap.-Nov. rainfall.
Mean of four years x five pasture treatments						
10	8040	4610	57	45	5.6	7.5
11	8660	4650	54	49	5.7	8.2
13	8170	5030	62	55	6.7	9.2
14	8620	5790	67	57	6.6	9.5
15	8550	5720	67	61	7.1	10.2
17	8580	6240	73	68	7.9	11.4

APPENDICES

Appendix Table 1

METEOROLOGICAL RECORDS FROM THE KANGAROO ISLAND RESEARCH CENTRE, PARNDANA EAST

1970

Climatic Data	MONTHLY MEANS												MEAN
	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.	
Cloud Cover [†]	0.60	0.57	0.61	0.65	0.55	0.53	0.58	0.70	0.80	0.68	0.60	0.61	0.62
Wind Speed (km hr ⁻¹)	22.1	22.6	19.4	18.4	16.6	19.2	17.0	20.6	20.5	22.6	21.0	24.5	20.4
Sea level Pressure (mb)	1015.4	1017.5	1017.1	1016.9	1022.4	1015.3	1019.3	1017.8	1015.4	1016.3	1016.2	1013.6	1016.9
Dry Bulb Temp. (°C)	17.7	19.3	16.8	15.8	11.7	11.2	10.1	9.5	10.3	13.4	16.4	18.5	14.2
Dew Pt. (°C)	10.9	11.8	10.8	11.5	9.4	9.0	8.1	6.9	7.2	8.1	8.8	9.8	9.3
Max. Temp (°C)	23.1	25.9	22.5	20.9	15.9	14.7	14.3	12.8	13.7	17.5	21.0	23.4	18.8
Min Temp. (°C)	11.3	12.9	10.9	11.1	7.9	8.0	7.1	5.8	6.5	6.6	8.3	10.9	8.9
Rainfall (mm) [‡]	40.3	2.8	21.7	51.0	49.6	93.0	49.6	111.6	81.0	18.6	27.0	27.9	570.7 ^S
No. of Wet Days	12	1	11	14	19	24	21	28	21	16	11	10	188 ^S
Sunshine (hrs day ⁻¹)	8.1	8.9	6.7	4.9	3.8	3.3	4.5	3.5	3.0	7.2	8.1	7.3	5.8

[†] 0 = no cloud cover 1 = maximum cloud cover

[‡] monthly totals. S = annual total.

Appendix Table 2

METEOROLOGICAL RECORDS FROM THE KANGAROO ISLAND RESEARCH CENTRE, PARNDANA EAST

1971

Climatic Data	MONTHLY MEANS												MEAN
	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.	
Cloud Cover [†]	0.61	0.64	0.63	0.60	0.66	0.74	0.64	0.57	0.59	0.64	0.75	0.70	0.65
Wind Speed (km hr ⁻¹)	21.0	21.7	22.6	18.9	15.9	14.7	14.8	18.7	19.7	23.8	20.6	24.5	19.8
Sea Level Pressure (mb)	1015.1	1013.3	1017.4	1017.4	1014.7	1020.8	1019.9	1014.4	1013.9	1011.5	1011.9	1011.6	1015.1
Dry Bulb Temp (°C)	19.0	19.5	19.7	16.6	12.4	10.8	9.8	9.9	11.8	13.3	14.3	16.4	14.5
Dew Pt. (°C)	12.9	12.9	14.1	12.4	10.2	9.0	8.1	7.6	8.8	8.0	9.4	11.2	10.4
Max Temp (°C)	24.6	26.0	24.4	22.2	15.8	14.2	13.2	13.5	14.9	17.0	17.7	20.5	18.7
Min Temp (°C)	12.4	13.6	14.6	12.0	9.1	7.7	6.5	6.3	7.0	7.5	9.4	10.2	9.7
Rainfall (mm) [‡]	6.2	11.2	34.1	138.0	96.1	66.0	62.0	142.6	66.0	37.2	81.0	62.0	791.5 ^S
No. of Wet Days	5	6	10	14	26	24	18	29	22	21	16	12	203 ^S
Sunshine (hr day ⁻¹)	7.7	6.9	5.9	5.9	3.9	2.5	3.0	4.4	5.2	5.0	4.6	5.9	5.1

[†] 0 = no cloud cover 1 = maximum cloud cover

[‡] monthly totals S = annual total.

Appendix Table 3

METEOROLOGICAL RECORDS FROM THE KANGAROO ISLAND RESEARCH CENTRE, PARNDANA EAST

1972

Climatic Data	MONTHLY MEANS												MEAN
	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.	
Cloud Cover ⁺	0.64	0.54	0.51	0.52	0.56	0.50	0.66	0.55	0.58	0.63	0.61	0.60	0.57
Wind Speed (km hr ⁻¹)	18.8	19.2	19.1	18.1	13.9	15.4	17.8	16.6	19.7	21.2	18.2	21.1	18.3
Sea Level Pressure (mb)	1013.9	1016.3	1018.5	1021.1	1027.9	1022.9	1017.1	1013.6	1020.2	1019.1	1018.3	1016.4	1018.8
Dry Bulb Temp (°C)	18.1	20.4	17.6	16.2	14.0	9.5	10.0	10.8	12.6	13.3	14.1	18.2	14.6
Dew Pt. (°C)	12.6	12.2	9.8	10.9	8.8	6.8	7.8	8.0	8.0	8.9	9.3	10.1	9.4
Max Temp (°C)	22.0	25.6	22.6	21.5	19.2	15.4	13.4	13.9	16.6	17.8	19.6	25.5	19.4
Min Temp (°C)	12.8	13.4	11.5	10.4	8.8	5.2	7.5	7.1	7.6	8.0	8.3	11.5	9.3
Rainfall (mm) †	58.9	49.3	6.2	24.0	21.7	63.0	148.8	124.0	18.0	40.3	12.0	6.2	571.0 ^S
No. of Wet Days	8	9	5	10	10	18	28	22	10	14	5	4	143 ^S
Sunshine (hr day ⁻¹)	6.8	7.1	6.2	6.1	4.9	4.5	2.6	4.4	5.1	6.2	6.5	8.2	5.7

⁺ 0 = no cloud cover 1 = maximum cloud cover

† monthly totals S = annual total.

METEOROLOGICAL RECORDS FROM THE KANGAROO ISLAND RESEARCH CENTRE, PARNDANA EAST

1973

Climatic Data	MONTHLY MEANS												MEAN
	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.	
Cloud Cover ⁺	0.67	0.44	0.63	0.59	0.45	0.59	0.37	0.59	0.56	0.63	0.59	0.52	0.55
Wind Speed(km hr ⁻¹)	19.4	17.2	17.2	21.3	19.2	15.5	15.3	18.9	26.2	21.2	14.6	21.3	18.9
Sea Level Pressure (mb)	1014.0	1015.7	1018.8	1017.0	1018.5	1017.6	1021.5	1016.3	1015.2	1017.0	1015.2	1013.3	1016.7
Dry Bulb Temp.(°C)	19.5	18.4	17.1	16.2	13.3	9.2	10.3	11.1	12.7	14.7	14.6	17.6	14.5
Dew Pt. (°C)	12.5	13.1	11.4	11.4	10.9	7.2	8.5	8.9	9.2	10.6	10.2	11.7	10.4
Max.Temp. (°C)	26.8	25.7	22.4	20.7	17.3	13.5	14.3	14.5	16.4	18.5	19.7	24.3	19.5
Min.Temp (°C)	14.6	12.4	11.8	11.4	8.9	6.3	6.5	7.2	8.0	8.5	8.7	11.6	9.7
Rainfall (mm)†	6.2	8.4	71.3	42.0	62.0	108.0	89.9	139.5	72.0	55.8	18.0	43.4	699.3 ^S
No. of Wet Days	9	7	11	15	15	17	16	24	20	16	16	12	178 ^S
Sunshine (hr day ⁻¹)	6.2	7.9	5.0	5.5	5.7	3.7	5.3	4.6	4.9	5.2	6.8	6.8	5.6

⁺0 = no cloud cover 1 = maximum cloud cover

† monthly totals S = annual total.

Appendix Table 5

METEOROLOGICAL RECORDS FROM THE KANGAROO ISLAND RESEARCH CENTRE, PARNDANA EAST

1974

Climatic Data	MONTHLY MEANS												MEAN
	JAN.	FEB.	MAR.	APR.	MAY	JUNE	JULY	AUG.	SEP.	OCT.	NOV.	DEC.	
Cloud Cover [†]	0.53	0.54	0.66	0.71	0.62	0.54	0.61	0.47	0.72	0.63	0.64	0.54	0.60
Wind Speed (km hr ⁻¹)	24.5	20.9	22.3	18.0	13.1	15.6	19.9	19.6	22.4	22.0	19.9	20.5	19.9
Sea Level Pressure (mb)	1010.9	1016.1	1017.5	1017.3	1020.5	1021.2	1013.0	1016.4	1018.5	1014.6	1015.9	1012.7	1016.2
Dry Bulb Temp. (°C)	20.0	17.2	19.6	15.2	12.7	11.0	10.5	10.9	11.4	13.5	13.9	16.4	14.4
Dew Pt. (°C)	14.7	13.9	14.9	12.8	11.2	9.7	8.2	9.2	8.1	10.0	9.7	10.1	11.0
Max. Temp. (°C)	26.2	23.9	25.3	19.0	16.1	14.6	13.5	14.7	14.5	17.3	18.2	22.1	18.8
Min. Temp. (°C)	14.9	13.5	15.1	11.8	9.8	7.7	7.5	7.3	6.2	8.7	7.9	9.9	10.0
Rainfall (mm) [‡]	49.6	8.4	24.8	60.0	108.5	66.0	161.2	80.6	72.0	93.0	27.0	9.3	761.7 ^S
No. of Wet Days	9	7	9	19	17	17	28	23	22	16	12	10	189 ^S
Sunshine (hr day ⁻¹)	7.2	7.5	4.8	1.7	3.2	3.4	2.7	5.2	4.0	5.5	9.0	9.2	5.3

[†]0 = no cloud cover 1 = maximum cloud cover

[‡]monthly totals S = annual total.

Appendix Table 6

MEAN PASTURE PRODUCTION MEASURED IN SEVEN PERIODS IN 1971

(kg ha⁻¹ dry matter)

Period	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)							Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means on SR.	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17						
24/4-2/6	2350	1035	2205	1710	2090	2030	1730	1795	1820	2030	1860	1875	***	200	NS	NS	
2/6-5/7	675	1085	905	1235	755	815	935	1040	1155	900	745	930	**	321	NS	NS	
5/7-8/9	4105	1490	2455	1645	2635	2210	2255	2365	2575	2650	2730	2640	***	506	*	NS	
8/9-24/11	3760	3955	3285	2645	4850	3945	4110	3145	3805	3840	3350	3700	***	751	NS	NS	
24/4-5/7	3025 ⁺	2120	3115	2945	2845	2840	2665	2840	2975	2930	2605	2810	-	-	NS	*	
24/4-8/9	7125 [*]	3610	5565	4590	5480	5050	4925	5205	5550	5580	5335	5270	***	724	NS	NS	
24/4-24/11	10885	7570	8850	7235	10325	9000	9035	8350	9350	9420	8685	8970	***	1236	NS	NS	

⁺ = Significance of within-treatment regressions on stocking rate.

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 7

MEAN PASTURE PRODUCTION MEASURED IN FIVE PERIODS IN 1972

(kg ha⁻¹ dry matter)

Period	Pasture treatments means				Stocking rate means (sheep ha ⁻¹)						Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means on SR.	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15						17
1/5-27/6	1365	590	270 [†]	500 [†]	835	795	760	755	655	565	745	710	***	427	NS	NS
27/6-26/9	2885	2925	1595	1680	2340	2225	2310	2320	2415	2070	2375	2280	***	464	NS	NS
26/9-16/11	2980	3355	3705	3485	3065	2930	3940	3445	3090	2985	3520	3320	NS	846	NS	NS
1/5-26/9	4250	3515	1870	2180	3190	3015	3070	3075	3065	2640	3140	3000	***	707	NS	NS
1/5-16/11	7230	6875	5575	5665	6240	5950	7010	6515	6155	5625	6640	6320	*	1178	NS	NS

† = Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 8
 MEAN PASTURE PRODUCTION MEASURED IN FIVE PERIODS IN 1973
 (kg ha^{-1} dry matter)

Period	Pasture treatment means					Stocking rate means (sheep ha^{-1})					Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. on SR	Hetero. Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15						17
18/3-18/6	2925	2700	2400	2610	2920	2580	2580	2805	2795	2620	2890	2710	NS	454	NS	NS
18/6-17/9	2615	2195 [†]	2605	3115	2730	2535	3355	2200	2940	2405	2480	2650	-	-	NS	*
17/9-26/11	3760 ^{**}	4295	3225	4365	3235	3105	3295	3400	3875	4405	4585	3780	*	929	***	NS
18/3-17/9	5540	4895	5010	5725	5650	5110	5935	5000	5735	5025	5370	5360	NS	1002	NS	NS
18/3-26/11	9300 [*]	9190	8230	10090	8885	8215	9235	8400	9610	9435	9955	9140	-	-	*	*

[†] = Significance of within-treatment regressions on stocking rate.

NS = $P > 0.05$ * = $P < 0.05$ ** = $P < 0.01$ *** = $P < 0.001$

Appendix Table 9
MEAN PASTURE PRODUCTION MEASURED IN FIVE PERIODS IN 1974

(kg ha⁻¹ dry matter)

Period	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)							Grand mean	Hetero. Adj't. treat. means	LSD 5% treat. means	Sig. mean Reg. Coeff. on SR.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17						
6/4-10/6	2880	3080 [†]	2775	2655	2490	2690	2830	2665	2950	2725	2795	2780	-	-	NS	**	
10/6-2/9	2130	2130 [*]	2210	2565	2585	1755	1870	2835	2435	2610	2435	2320	NS	1151	NS	NS	
2/9-18/11	4295	3970	4270 [*]	4065	4480	4565	4645	3925	3970	4370	3820	4220	NS	864	NS	NS	
6/4 -2/9	5010	5210	4985	5215	5075	4445	4700	5500	5385	5335	5230	5100	NS	1087	NS	NS	
6/4-18/11	9305	9180	9250	9280	9555	9010	9350	9425	9355	9705	9055	9320	NS	1081	NS	NS	

[†] = Significance of within-treatment regressions on stocking rate.

NS = P = > 0.05 * = P < 0.05 ** = P < 0.01 ***=P < 0.001

Appendix Table 10

MEAN CRUDE UTILIZATION OF PASTURE MEASURED IN SEVEN PERIODS IN 1971

(kg ha⁻¹ dry matter)

Period	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)							Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17					
24/4-2/6	1320 ^{*†}	845	1390	1210	1255	1220	870	1280	1155	1405	1285	1200	NS	443	NS	NS
2/6-5/7	625 [*]	590	710	680	755	365	565	700	930	715	750	670	NS	258	**	NS
5/7-8/9	3790	1930	3000	2440	2725	2575	2675	2500	2800	3115	2985	2775	***	421	*	NS
8/9-24/11	1155	3060	2230	2370	2885	2310	2365	1880	2560	2210	2720	2340	**	855	NS	NS
24/4-5/7	1945 [*]	1435	2100	1890	2005	1590	1435	1980	2085	2120	2035	1875	NS	494	**	NS
24/4-8/9	5730 [*]	3365	5100	4325	4730	4165	4110	4480	4890	5235	5020	4650	***	767	**	NS
24/4-24/11	6885	6425	7330	6700	7610	6475	6480	6360	7445	7440	7740	7000	NS	137	*	NS

+ = Significance of within-treatment regressions on stocking rate.

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 11

MEAN CRUDE UTILIZATION OF PASTURE MEASURED IN FIVE PERIODS IN 1972

(kg ha⁻¹ dry matter)

Period	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)					Grand mean	Hetero, Adj. treat, means	LSD 5% treat, means on SR.	Sig. mean Reg, Coeff.	Hetero, Reg. Coeff.	
	A	B	C	D	E	10	11	13	14	15						17
1/5-27/6	1255	360	150	415	610	500	555	620	550	460	665	560	-	-	NS	*
27/6-26/9	1940	2365	1405	1460 ⁺	2005	1980	1545	1895	1850	1675	2060	1830	**	522	NS	NS
26/9-16/11	1715	1295	2265 ^{**}	2120	1465 [*]	1480	1085	1580	1955	2005	2520	1770	*	605	***	NS
1/5-26/9	3195	2720	1555	1875 ^{**}	2610	2475	2100	2515	2395	2140	2725	2390	***	739	NS	NS
1/5-16/11	4910	4015	3820 ^{**}	3995	4075 [*]	3955	3185	4095	4355	4145	5245	4160	NS	1042	**	NS

+ = Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 12

MEAN CRUDE UTILIZATION OF PASTURE MEASURED IN FIVE PERIOD IN 1973
(kg ha⁻¹ dry matter)

Period	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)					Grand mean	Hetero. Adj. treat. means	Hetero. LSD 5% treat. means	Sig. mean Reg. on SR.	Hetero. Reg. Coeff.		
	A	B	C	D	E	10	11	13	14						15	17
18/3-18/6	1750 ^{*†}	2100 [*]	1940	2275	2220	1705	1675	2030	2175	2155	2605	440	NS	486	***	NS
18/6-17/9	1390	505 [*]	1575	2020	1895	885	1340	1230	1955	1605	1855	1480	-	-	*	**
17/9-26/11	2475	1970	1120	2015	575	1295	1940	1240	1815	1585	1915	1630	**	983	NS	NS
18/3-17/9	3145 [*]	2605	3520	4295	4110 ^{*†}	2590	3010	3260	4125	3765	4455	3530	*	1007	***	NS
18/3-26/11	5620 [*]	4575	4640	6310	4685	3885	4955	4495	5940	5345	6370	5160	NS	1425	**	NS

[†] = Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 13

MEAN CRUDE UTILIZATION OF PASTURE MEASURED IN FIVE PERIODS IN 1974

(kg ha⁻¹ dry matter)

Period	Pasture treatment means				Stocking rate means (sheep ha ⁻¹)					Grand mean	Hetero. Adj't. treat. means	LSD 5% treat. means on SR.	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.		
	A	B	C	D	E	10	11	13	14						15	17
6/4-10/6	1000	1615	1640	1250	1280	1035	930	1285	1500	1730	1665	1360	NS	579	**	NS
10/6-2/9	2045	1830 ⁺	1560	2240	2465	1965	1930	2270	2070	2210	1715	2030	NS	988	NS	NS
2/9-18/11	1320	1210	1705 ^{**}	1510	2555	1140	1130	1615	1845	2005	2225	1660	***	502	***	NS
6/4-2/9	3045	3445 [*]	3200	3490	3740	3000	2860	3555	3570	3940	3380	3380	NS	1092	NS	NS
6/4-18/11	4365	4655	4905	5000	6295	4140	3990	5170	5415	5945	5605	5040	*	1163	**	NS

+ = Significance of within-treatment regressions on stocking rate.

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 14

AVAILABILITY OF PASTURE MEASURED ON SEVEN OCCASIONS IN 1971-1972

(kg ha⁻¹ dry matter)

Date	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)							Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17						
2/6/71	1030	195	820	500	835	805	860	520	665	625	575	675	***	182	***	NS	
5/7/71	1080	685	1015	1060	840	1255	1230	860	890	810	570	935	**	218	***	NS	
8/9/71	1395	250	465 ⁺	265	750	885	810	735	660	345	315	625	***	330	***	NS	
24/11/71	4000	1145	1520 ⁺	540	2715	2525	2555	1990	1905	1980	950	1985	***	691	***	NS	
27/6/72	105	230	120	90	225	295	200	135	105	105	85	155	NS	146	**	NS	
26/9/72	1055	795	310	310	560	540	970	560	670	500	400	605	***	232	*	NS	
16/11/72	2320	2860	1755	1670	2165	1990	3830	2425	1800	1480	1395	2155	NS	1163	**	NS	

+ = Significance of within-treatment regression on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 15
AVAILABILITY OF PASTURE MEASURED ON SIX OCCASIONS IN 1973-1974
(kg ha⁻¹ dry matter)

Date	Pasture treatment means					Stocking rate means (sheep ha ⁻¹)						Grand mean	Hetero, Adj. treat, means	LSD 5% treat, means	Sig. mean Reg. Coeff.	Hetero, Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17					
18/6/73	1170	600 ⁺	460	335	705 ^{***}	875	905	775	620	465	285	655	***	227	***	NS
17/9/73	2395 [*]	2290	1490 [*]	1430 [*]	1540 ^{***}	2520	2925	1745	1610	1260	915	1830	**	564	***	NS
26/11/73	3680	4615 [*]	3590	3780	4200 [*]	4330	4280	3905	3665	4085	3580	3975	-	-	*	*
10/6/74	1880	1465 [*]	1135	1400	1215	1655	1905	1360	1450	995	1130	1415	NS	636	*	NS
2/9/74	1965	1765	1780	1730	1335	1440	1840	1945	1815	1395	1850	1715	NS	669	NS	NS
18/11/74	4940 [*]	4525 [*]	4345 [*]	4280	3260	4865	5355	4255	3940	3760	3450	4270	***	650	***	NS

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

+ = Significance of within-treatment regressions on stocking rate.

Appendix Table 16

MEAN LIVE WEIGHTS ON EIGHT OCCASIONS IN 1970-1971
(kg)

Date	Pasture treatment means					Stocking rate means (kg ha ⁻¹)							Grand mean	Hetero. Adj't. treat. means	LSD 5% treat. means on SR,	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17						
28/10/70	46.3	45.9	46.1	46.1	46.4	46.5	45.4	46.8	45.8	46.5	46.0	46.2	NS	-	NS	NS	
10/2/71	39.9	31.8	41.3	41.5	42.2	40.9	41.1	40.9	39.9	41.1	40.7	40.8	***	1.26	NS	NS	
31/3/71	39.4	30.1	43.7	42.8	44.9	42.4	42.3	41.3	40.5	42.4	41.8	41.8	***	2.20	NS	NS	
3/5/71	38.0	27.3	41.4	39.7	42.3	40.7	39.1	39.1	37.9	39.9	39.4	39.3	***	1.95	*	NS	
23/6/71	44.6 [†]	28.8	44.2	39.9 [†]	45.1 [*]	44.7	43.3	42.6	41.4	40.7	40.7	42.2	***	1.56	***	NS	
20/7/71	46.6 ^{**}	29.2 ^{**}	45.5 ^{**}	40.6 ^{**}	47.7	47.5	44.9	43.9	43.2	41.7	41.0	43.7	***	1.75	***	NS	
24/8/71	52.0 [*]	30.1 [*]	47.0 ^{**}	41.2 ^{**}	51.2 [*]	51.3	48.2	46.7	45.8	43.3	42.1	46.2	***	2.40	***	NS	
5/10/71	53.1 ^{**}	30.1 ^{**}	46.7 [*]	39.7 ^{**}	51.0 ^{**}	49.6	50.0	45.9	44.9	43.0	43.6	46.2	***	2.48	***	NS	

[†] Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 17
 MEAN LIVE WEIGHTS ON SEVEN OCCASIONS IN 1971-1972
 (kg)

Date	Pasture treatment means				Stocking rate means (kg ha ⁻¹)							Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. Coeff. on SR.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17					
11/11/71	57.6	*** 38.1	53.6	*** 45.4	58.0	54.6	55.0	51.8	52.8	51.0	51.8	52.8	-	***	***	***
6/12/71	59.2	*** 41.9	55.9	*** 50.2	* 60.1	58.1	58.1	55.6	55.3	53.3	53.8	55.7	-	***	***	**
22/12/71	58.7	** 43.7	56.5	** 52.2	60.0	58.0	59.2	56.2	56.5	54.7	53.7	56.4	-	***	***	*
19/1/72	53.6	*** 41.7	54.4	*** 49.2	54.7	54.5	54.8	53.1	53.3	51.1	49.1	52.7	-	***	***	*
SHEARING January 25, 1972																
25/1/72	43.4	* 41.5	44.4	** 40.3	44.5	44.1	44.0	43.3	43.3	42.3	40.8	43.0	2.04	**	**	NS
8/3/72	49.1	44.4	49.9	** 44.6	* 49.7	50.0	49.6	47.8	47.4	47.5	43.8	47.7	2.44	***	***	NS
19/4/72	* 50.0	43.1	46.3	* 42.0	48.9	49.6	49.9	46.4	44.6	46.0	41.0	46.2	3.06	***	***	NS

+ Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 18

MEAN LIVE WEIGHTS ON SEVEN OCCASIONS IN 1972-1973

(kg)

Date	Pasture treatment means				Stocking rate means (kg ha ⁻¹)							Grand mean	Hetero. Adj't. treat. means	LSD 5% treat. means on SR.	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17					
25/5/72	47.5	** 40.9	** 43.0	** 38.9	* 46.1	47.1	47.7	43.2	42.4	42.1	38.9	43.6	**	2.65	**	NS
12/7/72	** 42.6	* 36.9	* 38.8	* 35.0	* 41.9	44.1	44.2	38.4	37.5	36.9	34.3	39.2	**	2.71	**	NS
16/8/72	* 46.3	* 40.3	** 39.4	* 37.7	** 43.4	45.3	46.2	41.2	40.0	40.4	36.5	41.6	**	2.39	**	NS
28/9/72	48.1	** 42.0	** 39.2	* 36.9	** 43.7	46.0	47.5	42.4	41.3	40.1	35.8	42.2	**	2.84	**	NS
29/11/72	58.4	** 56.3	* 54.6	* 50.1	56.0	57.4	57.9	56.3	54.8	54.8	50.3	55.3	-	-	**	*
17/1/73	59.1	60.1	58.0	* 52.3	58.5	59.9	60.3	59.5	57.5	58.5	50.7	57.7	-	-	**	*
18/1/73	50.1	51.1	49.0	* 44.7	49.2	51.1	50.3	50.1	48.2	49.3	44.0	48.8	*	3.87	**	NS

+ Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 19

MEAN LIVE WEIGHTS ON EIGHT OCCASIONS IN 1973-1974

(kg)

Date	Pasture treatment means					Stocking rate means (kg ha ⁻¹)							Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. onSR.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17						
28/2/73	53.9	56.1	52.9	49.4 ^{**†}	53.1	55.2	55.9	55.1	52.0	53.1	47.6	53.2	NS	4.39	***	NS	
18/4/73	54.0	55.1 ^{**}	48.8 [*]	47.4 ^{**}	51.6 ^{**}	55.5	56.1	53.6	50.1	49.6	43.6	51.4	-	-	***	*	
5/6/73	55.5	56.5 [*]	51.2 [*]	48.2 ^{**}	54.6 [*]	58.2	57.4	54.9	51.6	50.9	46.4	53.2	***	3.85	***	NS	
17/7/73	61.3	60.5 ^{**}	56.1 [*]	52.9 ^{**}	58.6 [*]	63.7	61.2	59.8	56.4	55.6	50.7	57.9	-	-	***	*	
21/8/73	64.5	64.2	59.8 [*]	56.9 ^{**}	62.5 [*]	66.2	64.4	62.3	60.7	59.7	56.3	61.6	-	-	***	*	
3/10/73	65.6	64.9	61.6	60.5 ^{**}	64.4 [*]	67.8	64.6	64.2	61.8	62.7	59.2	63.4	*	3.26	***	NS	
20/11/73	70.8	70.0	68.9	67.8	70.9	72.5	68.6	69.6	69.6	69.4	68.2	60.6	NS	2.83	NS	NS	
15/1/74	66.4	63.8	64.8	63.0 ^{**}	67.5	68.3	64.4	66.4	64.7	63.5	63.2	65.1	*	3.12	*	NS	

SHEARING January 15, 1974

[†]Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 20

MEAN LIVE WEIGHTS ON EIGHT OCCASIONS IN 1974-1975
(kg)

Date	Pasture treatment means				Stocking rate means (kg ha ⁻¹)					Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. on SR.	Hetero. Reg. Coeff.		
	A	B	C	D	E	10	11	13	14						15	17
5/3/74	58.0	55.9	58.0	57.7 ^{*†}	59.5	59.6	57.4	58.7	57.4	57.0	56.7	57.8	NS	2.86	NS	NS
1/5/74	53.9	54.1	53.4	53.8 ^{***}	54.3 ^{**}	57.7	56.2	53.7	52.9	52.4	50.8	54.0	NS	2.30	***	NS
6/6/74	57.6	58.0 [*]	56.1	57.3 ^{**}	57.1 [*]	60.8	58.8	57.4	56.1	55.4	54.6	57.2	NS	2.39	***	NS
18/7/74	63.6	62.5	61.6	62.7	61.6	64.7	63.1	62.4	63.2	60.1	60.8	62.4	NS	2.83	**	NS
11/9/74	67.1	65.7	64.7	66.4	65.4	67.8	65.1	65.8	66.2	64.7	65.2	65.8	NS	2.70	NS	NS
30/10/74	69.5	70.0	69.7	69.8	69.7	70.6	68.3	69.8	70.6	69.5	69.4	69.7	NS	3.11	NS	NS
21/11/74	69.6	70.5	69.8	70.3	70.9	71.4	68.5	70.8	69.9	70.3	70.0	70.2	NS	3.12	NS	NS
23/1/75	68.7	68.0	68.9	68.5	70.2	70.7	66.6	70.1	69.0	68.5	68.2	68.9	NS	3.02	NS	NS
SHEARING January 23, 1975																

[†] = Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 21

REGRESSION COEFFICIENTS OF LIVE WEIGHT ON STOCKING RATE ON 20
OCCASIONS DURING 1971-1973

Date	Pasture treatments				
	A	B	C	D	E
10/2/71	-0.20	-0.47	-0.04	+0.02	-0.16
31/3/71	-0.43	-0.70	-0.31	-0.19	+0.06
3/5/71	-0.18	-0.95	-0.50	-0.29	-0.12
23/6/71	-1.21	-0.40	-1.16	-0.80	-0.67
20/7/71	-1.42	-1.23	-1.41	-1.04	-1.12
24/8/71	-1.98	-1.59	-1.91	-1.47	-1.58
5/10/71	-1.20	-2.22	-1.56	-1.88	-1.46
11/11/71	-0.19	-2.06	-1.02	-1.75	-0.29
6/12/71	-0.34	-1.93	-1.23	-1.91	-0.50
22/12/71	-0.26	-1.87	-0.86	-1.73	-0.63
19/1/72	-0.54	-2.10	-1.01	-1.59	-0.17
	SHEARING				
25/1/72	-0.32	-0.62	-0.32	-1.00	-0.25
8/3/72	-1.07	-0.99	-0.59	-1.24	-0.37
19/4/72	-1.06	-1.63	-0.92	-1.99	-0.72
25/5/72	-1.03	-2.02	-1.26	-1.66	-0.72
12/7/72	-1.88	-2.18	-1.21	-1.62	-1.01
16/8/72	-1.49	-1.75	-1.20	-1.10	-1.42
28/9/72	-1.28	-2.18	-1.13	-1.79	-1.78
29/11/72	+0.04	-1.06	-0.91	-2.22	-0.94
17/1/73	+0.13	-0.83	-1.00	-2.89	-1.13
	SHEARING				

Appendix Table 22

REGRESSION COEFFICIENTS OF LIVE WEIGHT ON STOCKING RATE ON 17
OCCASIONS DURING 1973-1975

Date	Pasture treatments				
	A	B	C	D	E
18/1/73	-0.26	-0.36	-0.84	-2.09	-1.05
28/2/73	+0.15	-0.69	-1.80	-1.99	-0.97
18/4/73	-0.50	-1.50	-2.22	-2.80	-1.88
5/6/73	-0.80	-1.21	-2.16	-2.71	-1.94
17/7/73	-0.63	-1.10	-2.17	-2.96	-2.23
21/8/73	-0.52	-0.96	-1.62	-2.44	-1.58
3/10/73	-0.53	-0.41	-1.08	-2.01	-1.43
20/11/73	-0.22	-0.10	-0.65	-0.55	-0.49
15/1/74	-0.64	+0.15	-0.54	-1.08	-0.74
	SHEARING				
5/3/74	-0.20	+0.27	+0.20	-0.92	-0.89
1/5/74	-0.50	-0.60	-0.94	-1.47	-1.32
6/6/74	-0.38	-0.59	-0.85	-1.40	-1.15
18/7/74	-0.18	-0.75	-0.45	-1.07	-0.33
11/9/74	-0.11	-0.39	-0.27	-0.49	-0.13
30/10/74	+0.17	-0.36	-0.02	-0.12	+0.13
21/11/74	+0.01	-0.17	+0.12	-0.06	-0.08
23/1/75	-0.42	-0.23	+0.22	-0.14	-0.05

Appendix Table 23

MEAN GROWTH RATE OF CLEAN WOOL IN SEVEN PERIODS IN 1971-1972

(g day⁻¹)

Period	Pasture treatment means					Stocking rate means (kg ha ⁻¹)							Grand mean	Hetero. Adj't. treat. means	LSD 5% treat. means	Sig. mean Reg. on SR.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17						
28/1-31/3	9.7	9.0 [†]	10.8	9.1 ^{**}	9.8	10.5	10.4	9.8	9.7	9.7	9.2	9.7	NS	-	**	**	
31/3-3/5	12.4	9.6	12.5	9.5 [*]	15.0	12.4	14.0	12.2	11.9	11.6	11.6	11.8	-	-	**	*	
3/5-24/6	8.9	5.4	8.6 ^{**}	6.1 ^{**}	9.4	9.0	9.2	8.2	7.9	7.2	7.5	7.7	-	-	***	*	
24/6-25/8	11.4	6.1	10.0 ^{***}	8.0 ^{**}	12.1	11.7	11.4	10.5	10.0	9.1	8.2	9.5	***	1.15	***	NS	
25/8-5/10	13.1	6.7	12.7 ^{**}	9.6 [*]	14.0	13.2	13.3	12.7	11.9	10.1	11.1	11.2	***	1.52	***	NS	
5/10-11/11	14.7 [*]	10.8	17.0	14.6 [*]	18.1	17.0	17.7	15.9	15.9	14.1	13.9	15.1	***	2.2	**	NS	
11/11-28/1/72	4.5	3.5	6.3	4.3	5.3 [*]	5.4	5.3	4.9	4.9	4.3	3.4	4.8	NS	1.89	**	NS	

[†] Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 24

MEAN GROWTH RATE OF CLEAN WOOL IN EIGHT PERIODS IN 1972-1973

(g day⁻¹)

Period	Pasture treatment means				Stocking rate means (kg ha ⁻¹)				Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means on SR.	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.			
	A	B	C	D	E	10	11	13						14	15	17
28/1-8/3	13.1	13.2	13.7	12.4 ⁺	13.7	14.1	14.1	13.8	13.0	13.2	11.0	13.2	***	1.70	***	NS
8/3-19/4	16.1	15.6 ^{***}	15.1 ^{**}	14.0 [*]	15.9	16.6	16.7	15.9	14.8	14.8	13.2	15.3	**	1.03	***	NS
19/4-25/5	13.0	11.9 ^{***}	10.3 [*]	10.0	11.6 ^{**}	12.3	13.4	11.9	11.1	10.3	9.0	11.3	***	1.23	***	NS
25/5-12/7	7.2 ^{**}	6.2 ^{**}	6.0	5.6	6.8 [*]	7.3	7.2	6.8	5.8	6.3	4.9	6.4	**	0.73	***	NS
12/7-16/8	12.0 [*]	9.7 ^{**}	9.6 ^{**}	8.3	10.6 ^{**}	12.5	12.5	9.7	9.5	9.1	7.4	10.0	***	1.23	***	NS
16/8-28/9	7.8 [*]	6.5 [*]	6.5 [*]	5.3	6.9 ^{**}	7.8	8.1	6.4	6.4	6.1	5.2	6.6	***	0.92	***	NS
28/9-29/11	6.7	5.4	5.3	4.7	5.5	6.5	6.6	5.5	5.6	4.9	4.4	5.5	***	0.66	***	NS
29/11-18/1/73	10.8	9.8	10.2	10.1	10.0	11.3	11.5	10.2	9.8	9.8	8.4	10.2	NS	1.28	***	NS

⁺ = Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 25

MEAN GROWTH RATE OF CLEAN WOOL IN EIGHT PERIODS IN 1973-1974

(g day⁻¹)

Period	Pasture treatment means					Stocking rate means (kg ha ⁻¹)							Grand mean	Hetero. Adj. treat. means	LSD 5% treat. means	Sig. mean Reg. Coeff. on SR.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17						
18/1-28/2	16.3	15.6	16.7 ^{***†}	15.7	16.8	17.3	16.1	16.6	16.2	15.5	15.7	16.2	NS	1.33	*	NS	
28/2-18/4	16.1	16.4	16.8	16.1	16.5	16.5	15.6	16.2	16.3	16.4	17.1	16.4	NS	1.10	NS	NS	
18/4-5/6	15.9	16.3	16.1	15.6	16.0	16.4	15.7	15.7	15.7	16.0	16.5	16.0	NS	1.03	NS	NS	
5/6-17/7	14.1	14.1	14.1	12.8	13.6	14.4	13.9	13.3	13.8	13.4	13.6	13.7	NS	1.04	NS	NS	
17/7-21/8	21.4	20.6	20.4	18.4	20.3	22.1	20.9	19.9	19.7	19.9	18.8	20.2	-	-	***	*	
21/8-3/10	18.2 ^{**}	18.1	16.2 ^{**}	15.3 [*]	17.9	19.9	18.5	17.3	16.8	15.8	14.5	17.2	**	1.68	***	NS	
3/10-20/11	13.6	14.3	12.4 [*]	11.4 [*]	12.5 [*]	13.5	14.3	13.2	12.7	12.4	11.0	12.8	**	1.47	***	NS	
20/11-15/1/74	11.6	12.0	12.9	11.1	12.4	12.0	12.9	12.7	11.4	12.4	10.7	12.0	NS	1.25	*	NS	

† = Significance of within-treatment regressions on stocking rate

NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.001

Appendix Table 26

MEAN GROWTH RATE OF CLEAN WOOL IN SEVEN PERIODS IN 1974-1975
(g day⁻¹)

Period	Pasture treatment means					Stocking rate means (kg ha ⁻¹)							Grand mean	Hetero Adj. treat. means	LSD 5% treat. means on SR.	Sig. mean Reg. Coeff.	Hetero. Reg. Coeff.
	A	B	C	D	E	10	11	13	14	15	17						
15/1-1/5	6.0	6.0	6.3	6.0	6.3	6.3	5.8	6.6	5.9	6.1	6.0	6.1	NS	0.64	NS	NS	
1/5-6/6	14.1	14.7	16.3	14.2	16.4	14.3	15.6	16.5	14.1	14.2	15.8	15.1	NS	2.81	NS	NS	
6/6-18/7	17.7	17.4	17.7	17.2	18.4	17.7	17.2	17.5	17.7	17.6	18.3	17.7	NS	1.17	NS	NS	
18/7-11/9	13.1	13.0	12.5	13.0	13.0	12.7	12.7	12.4	13.2	13.1	13.5	12.9	NS	0.89	*	NS	
11/9-30/10	11.9	11.3	11.3	11.3	11.0 [†]	11.7	11.5	10.8	11.4	11.2	11.6	11.4	NS	1.06	NS	NS	
30/10-21/11	22.4	22.1	22.2	22.1	21.3	24.3	22.9	22.7	20.6	20.6	20.9	22.0	NS	2.54	**	NS	
21/11-23/1/75	13.2	13.5	14.4	15.3	14.0	15.7	14.8	14.4	13.4	13.7	12.5	14.1	NS	1.49	***	NS	

† = Significance of within-treatment regressions on stocking rate
 NS = P > 0.05 * = P < 0.05 ** = P < 0.01 *** = P < 0.01

Appendix Table 27

REGRESSION COEFFICIENTS OF CLEAN WOOL GROWTH RATE ON STOCKING RATE
IN 15 PERIODS (1971-1973)

Period	Pasture treatments				
	A	B	C	D	E
	<u>1971/72</u>				
28/1-31/3	-0.17	-0.60	-0.11	-0.65	+0.29
31/3-3/5	-0.03	-0.70	-0.45	-1.01	+0.14
3/5-24/6	-0.24	-0.53	-0.37	-0.60	-0.13
24/6-25/8	-0.55	-0.64	-0.54	-0.81	-0.45
25/8-5/10	-0.47	-0.19	-0.67	-0.66	-0.33
5/10-11/11	-0.54	-0.41	-0.32	-1.04	-0.17
11/11-28/1/72	-0.22	-0.03	-0.97	-0.17	-0.32
	<u>1972/73</u>				
28/1-8/3	-0.11	-0.44	-0.31	-0.96	-0.20
8/3-19/4	-0.35	-0.51	-0.49	-0.81	-0.35
19/4-25/5	-0.33	-0.81	-0.37	-0.55	-0.73
25/5-12/7	-0.45	-0.26	-0.24	-0.36	-0.36
12/7-16/8	-0.79	-0.93	-0.72	-0.64	-0.87
16/8-28/9	-0.39	-0.55	-0.45	-0.42	-0.31
28/9-29/11	-0.45	-0.47	-0.21	-0.39	-0.25
29/11-18/1/73	-0.17	-0.37	-0.43	-0.52	-0.64

Appendix Table 28

REGRESSION COEFFICIENTS OF CLEAN WOOL GROWTH RATE ON STOCKING RATE
ON 15 OCCASIONS (1973-75)

Period	Pasture treatments				
	A	B	C	D	E
	<u>1973/74</u>				
18/1-28/2	-0.06	+0.03	-0.49	-0.43	+0.02
28/2-18/4	+0.17	+0.28	-0.15	+0.04	+0.24
18/4-5/6	+0.12	+0.08	-0.26	-0.08	+0.28
5/6-17/7	+0.09	+0.03	-0.40	-0.30	+0.01
17/7-21/8	+0.13	-0.01	-1.06	-0.72	-0.41
21/8-3/10	-0.36	-0.60	-1.26	-0.90	-0.56
3/10-20/11	+0.12	-0.40	-0.62	-0.67	-0.46
20/11-15/1/74	-0.15	+0.10	-0.38	-0.37	-0.20
	<u>1974/75</u>				
15/1-1/5	-0.09	+0.03	-0.10	+0.05	0.00
1/5-6/6	+0.43	-0.05	-0.59	+0.49	-0.09
6/6-18/7	+0.25	+0.12	-0.29	+0.16	+0.22
18/7-11/9	+0.20	+0.13	-0.08	+0.04	+0.32
11/9-30/10	-0.01	0.00	-0.21	-0.06	+0.20
30/10-21/11	-0.34	-0.39	-0.91	-0.74	-0.14
21/11-23/1/75	-0.35	-0.29	-0.36	-0.52	-0.50

Appendix Table 29

NUMBER AND CAUSE OF DEATHS OF EXPERIMENTAL WETHERS 1970-1974.

Date	Cause of Death	No. Deaths	Treatment ⁺	Liveweight range at Death (kg)
<u>1971</u>				
April	Unknown	1	B11	26.3
May	Undernutrition	3	2xB17, B15	28.7 - 33.4
May	Unknown	1	C17	40.0
June	Undernutrition	3	2xB17, B14	27.6 - 35.0
July	Undernutrition	2	B17, B15	28.3 - 28.6
August	Undernutrition	1	B15	27.6
September	Unknown	1	B14	31.8
September	Undernutrition	2	2xD17	28.6 - 31.5
October	Undernutrition	1	D14	30.4
December	Flystrike	1	E10	49.6
Annual Total		16		
<u>1972</u>				
March	Shearing wounds	2	A17, C17	34.2 - 41.6
April	Undernutrition	2	D17, D15	32.4 - 37.2
May	Ruptured bladder	1	B17	44.5
July	Undernutrition	5	C17, B17, B15, D15, D14	24.4 - 36.0
June	Unknown	1	D14	
Annual Total		11		
<u>1973</u>				
January	Heat exhaustion	2	B11, A10	54.2 - 61.7
April	Unknown	1	D11	59.1
May	Unknown	1	E13	52.0
October	Unknown	1	D17	50.7
December	Urinary infection	1	A10	62.2
Annual Total		6		
<u>1974</u>				
January	Unknown	1	B10	65.0
January	Shearing wounds	3	B10, B14, C14	56.4 - 63.6
February	Unknown	1	D15	60.8
August	Unknown	1	A11	45.8
Annual Total		6		

⁺ = A, B, C, D and E indicate pasture treatments and postscripts indicate stocking rate (sheep ha⁻¹).

BIBLIOGRAPHY

11. BIBLIOGRAPHY

- ALLDEN, W.G. (1969). - The summer nutrition of weaner sheep: The voluntary feed intake, body weight change, and wool production of sheep grazing the mature herbage of sown pasture in relation to the intake of dietary energy under a supplementary feeding regime.
Aust. J. Agric. Res. 20: 499-512.
- ANON (1962). - Phalaris hybrids promising.
Rural Res. CSIRO. 41: 16-21.
- ARMSTRONG, D.G., BLAXTER, K.L. and WAITE, R. (1964). - The evaluation of artificially dried grass as a source of energy for sheep. III. The prediction of nutritive value from chemical and biological measurements.
J. Agric. Sci. 62: 417-424.
- ARNOLD, G.W. (1962). - Effects of pasture maturity on the diet of sheep.
Aust. J. Agric. Res. 13: 701-706.
- ARNOLD, G.W. (1964). - Some principles in the investigation of selective grazing.
Proc. Aust. Soc. Anim. Prod. 5: 258-271.
- ARNOLD, G.W. and McMANUS, W.R. (1960). - The effect of level of stocking on two pasture types upon wool production and quality.
Proc. Aust. Soc. Anim. Prod. 3: 63-68.
- AXELSEN, A. and MORLEY, F. (1968). - Evaluation of eight pastures by animal production.
Proc. Aust. Soc. Anim. Prod. 7: 92-98.
- BARCLAY, P.C. (1963). - Ariki, a new selected perennial type ryegrass variety.
Proc. N.Z. Grassl. Assoc. 25: 128-149.
- BARNARD, C. (1972). - Reg. Aust. Herb. Pl. cvv. CSIRO. (Aust.), pp. 16-17.
- BARRETT, D.W., CAMPBELL, N.A. and ARNOLD, G.W. (1973). - The effects of pre-maturity desiccation with paraquat and leaching on the quality of dry Wimmera ryegrass.
Aust. J. Exp. Agric. Anim. Husb. 13: 411-417.
- BARROW, N.J. (1969). - Implications of stocking rate experiments - nutrient requirement and fertilizer requirement.
Proc. Aust. Grassl. Conf., Perth, 1968. Vol. 2, pp. 169-171.
- BARTON, R.A. and ULYATT, M.J. (1963). - Pasture type in relation to live-weight gain, carcass composition, iodine nutrition and some rumen characteristics of sheep. II. Carcass composition and the non-carcass components of liveweight.
J. Agric. Sci. Camb. 61: 191-195.

- BATES, G.H. (1935). - The vegetation of footpaths, sidewalks, cart-tracks and gateways.
J. Ecol. 23: 470-487.
- BEALE, P.E. (1972). - Clover wilt threat to pastures.
J. Dep. Agric. S. Aust. 75: 69-71.
- BEALE, P.E. (1974). - Regeneration of *Trifolium subterraneum* cv. Yarloop from seed reserves in soil on Kangaroo Island.
J. Aust. Inst. Agric. Sci. 40: 78-80.
- BEALE, P.E., CHAMBERLAIN, H.V., NINNES, B.A. and LEWIS, M.R. (1976). - Trikkala - a replacement for Yarloop.
Dep. Agric. Fish. Spec. Bull. No. 2.76.
- BECK, A.B. (1952). - The composition of the Dwalganup strain of subterranean clover.
J. Agric. W. Aust. (3) 1: 257-259.
- BENNETTS, H.W., UNDERWOOD, E.J. and SHIER, F.L. (1946). - A specific breeding problem of sheep associated with subterranean clover pastures.
Aust. Vet. J. 22: 2-12.
- BIDDISCOMBE, E.F. (1953). - A survey of the natural pastures of the Trangie district, New South Wales, with particular reference to the grazing factor.
Aust. J. Agric. Res. 4: 1-28.
- BROWN, K.R. (1968). - The influence of herbage height at treading and treading intensity on the yield and botanical composition of a perennial ryegrass - white clover pasture.
N.Z. J. Agric. Res. 11: 131-137.
- BROWN, T.H. (1976a). - Effect of deferred autumn grazing and stocking rate of sheep on pasture production in a Mediterranean-type climate.
Aust. J. Exp. Agric. Anim. Husb. 16: 181-188.
- BROWN, T.H. (1976b). - The effect of stocking rate and deferred autumn grazing of pasture on liveweight and wool production of Merino wethers in a Mediterranean-type climate.
Aust. J. Exp. Agric. Anim. Husb. 16: 189-196.
- BROWN, T.H. (1977). - Rate of loss of dry matter and change in chemical composition of nine pasture species over summer.
Aust. J. Exp. Agric. Anim. Husb. 17: 75-79.
- BUNTON, G.W. and ANDREWS, J.S. (1948). - Recovery and viability of seeds of certain Southern Grasses and Lespedeza through the bovine digestive tract.
J. Agric. Res. 76: 95-103.
- CADE, J.W. (1969). - The ryegrasses in Victoria.
J. Dep. Agric. Vict. 67: 300-305.

- CAMERON, I.H. (1966). - Dry matter and quality changes during and after making pasture hay.
Aust. J. Exp. Agric. Anim. Husb. 6: 394-401.
- CAMERON, I.H. and CANNON, D.J. (1970). - Changes in the botanical composition of pasture in relation to rate of stocking with sheep, and consequent effects on wool production.
Proc. XI. Int. Grassl. Cong. Qld. Aust. pp. 640-643.
- CAMPBELL, R.J. and BEALE, J.A. (1973). - Evaluation of natural pastures at Trangie in central Western New South Wales.
2. Botanical composition changes with particular reference to *Hordeum leporinum*.
Aust. J. Exp. Agric. Anim. Husb. 13: 662-668.
- CANNON, D.J. (1969). - The wool production and liveweight of wethers in relation to stocking rate and superphosphate application.
Aust. J. Exp. Agric. Anim. Husb. 9: 172-180.
- CANNON, D.J. (1972). - The influence of rate of stocking and application of superphosphate on the production and quality of wool, and on the gross margin from Merino wethers.
Aust. J. Exp. Agric. Anim. Husb. 12: 348-354.
- CARTER, E.D. (1958). - Pasture renovation of the Stringybark Lands of Kangaroo Island.
J. Dept. Agric. S. Aust. 61: 401-408.
- CARTER, E.D. (1965). - Some relationships between superphosphate use and consequent animal production from pasture in South Australia.
Proc. IX. Int. Grassl. Cong., Sao Paulo, Brazil. Vol. 2, pp. 1027-1032.
- CARTER, E.D. (1968a). - The effect of stocking rate on the production and utilization of annual pasture and on wool production.
Proc. Aust. Grassl. Conf., Perth, 1968. Vol. 1, Sect. 1a, pp. 9-11.
- CARTER, E.D. (1968b). - Changes in seed supply of an annual pasture grazed continuously by sheep at five stocking rates.
Proc. Aust. Grassl. Conf., Perth, 1968. Vol. 1, Sect. 1b, pp. 7-10.
- CARTER, E.D. (1969). - Implications of stocking rate for botanical composition of pasture.
Proc. Aust. Grassl. Conf., Perth, 1968. Vol. 2, pp. 173-178.
- CARTER, E.D. (1970). - Phosphorous fertilizer requirements of the Seddon Soil Series on Kangaroo Island, South Australia.
M.Ag.Sc. Thesis, University of Adelaide, Adelaide, S. Aust.
- CARTER, E.D. (1973). - Pasture production and utilization.
Bi. Rep. Waite Agric. Res. Inst. 1972-73, pp. 42-43.

- CARTER, E.D., CHALLIS, S. and RIDGWAY, I.G. (1977). - The use of heavy solvents for separating seed from soil.
Proc. XIII. Int. Grassl. Cong., Leipzig, GDR. (in press).
- CARTER, E.D. and DAY, H.R. (1970). - Interrelationships of stocking rate and superphosphate rate on pasture as determinants of animal production.
Aust. J. Agric. Res. 21: 473-491.
- CARTER, E.D. and SIVALINGAM, T. (1977). - Some effects of treading by sheep on pastures of the Mediterranean climate zone of South Australia.
Proc. XIII. Int. Grassl. Cong., Leipzig, GDR. (in press).
- COCKS, P.S. (1974). - The influence of density and nitrogen on the outcome of competition between two annual pasture grasses (*Hordeum leporinum* Link and *Lolium rigidum* Gaud.).
Aust. J. Agric. Res. 25: 247-258.
- CONNIFFE, D., BROWNE, D. and WALSH, M.J. (1970). - Experimental design for grazing trials.
J. Agric. Sci. Camb. 74: 339-342.
- CONRAD, H.R. (1966). - Symposium on factors influencing the voluntary intake of herbage by ruminants: physiological and physical factors limiting feed intake.
J. Anim. Sci. 25: 227-235.
- COOK, L.J. (1939). - Further results in "topdressing" poor pasture lands with phosphatic fertilizers.
J. Dep. Agric. S. Aust. 42: 791-808 and 851-865.
- CRAWFORD, E.J. (1956). - Perennial grasses - Kangaroo Island Research Centre.
Dept. Agric. S. Aust. Pl. Introd. Sect. Ann. Rep. 1965/66, pp. 60-68.
- DAVIDSON, J.L. (1969). - Growth of grazed plants.
Proc. Aust. Grassl. Conf., Perth, 1968. Vol. 2, pp. 125-137.
- DAVIES, H.L. (1962a). - Studies of time of lambing in relation to stocking rate in south-western Australia.
Proc. Aust. Soc. Anim. Prod. 4: 113-120.
- DAVIES, H.L. (1962b). - Intake studies in sheep involving high fluid intake.
Proc. Aust. Soc. Anim. Prod. 4: 167-171.
- DAVIES, H.L. and GREENWOOD, E.A.N. (1972). - Liveweight and wool growth of sheep in response to the quantity and botanical composition of annual pasture induced by nitrogen fertilizer and stocking treatments.
Aust. J. Agric. Res. 23: 1101-1111.

- DAVIES, H.L. and HUMPHRIES, A.W. (1965). - Stocking rate and wool production at Kojonup. 1. Wether sheep.
J. Dep. Agric. W. Aust. 6: 409-413.
- DAVIES, W. (1938). - Vegetation of grass verges and other excessively trodden habitats.
J. Ecol. 26: 38-49.
- DAVIS, I.F. (1964). - Diet selected by sheep grazing on annual pasture in southern Victoria.
Proc. Aust. Soc. Anim. Prod. 5: 249-250.
- DAVIS, I.F. and SHARKEY, M.J. (1972). - Effect of fertilizer nitrogen and feeding practices on productivity of Corriedale wethers grazing annual pasture in southern Victoria.
Aust. J. Exp. Agric. Anim. Husb. 12: 589-595.
- DAY, H.R. (1963). - Sub-clovers for Kangaroo Island.
J. Dep. Agric. S. Aust. 66: 288-293.
- DAY, H.R. (1964). - Sub-clover strains under grazing.
Kangaroo Island Res. Centre Ann. Rep. 1963/64, pp. 9-10.
- DONALD, C.M. (1959). - The production and life span of subterranean clover (*Trifolium subterraneum* L.).
Aust. J. Agric. Res. 10: 771-787.
- DONALD, C.M. (1970). - Temperate pasture species. In "Australian Grasslands".
Ed. R.M. Moore (Aust. Nat. Uni. Press, Camb. Aust.)
pp. 303-320.
- DONALD, C.M. and ALLDEN, W.G. (1959). - The summer nutrition of weaner sheep: The deficiencies of the mature herbage of sown pasture as a feed for young sheep.
Aust. J. Agric. Res. 10: 199-218.
- DORE, W.G. and RAYMOND, L.C. (1942). - Viable seed in pasture soil and manure.
Sci. Agric. 23(2): 69-79.
- DRAKE, F.R. and ELLIOTT, N.M. (1963). - Rates of stocking investigated at Bengworden.
J. Dep. Agric. Vict. 61: 553-558.
- EDMOND, D.B. (1958). - Some effects of soil physical condition on ryegrass growth.
N.Z. J. Agric. Res. 1: 652-659.
- EDMOND, D.B. (1964). - Some effects of sheep treading on the growth of 10 pasture species.
N.Z. J. Agric. Res. 7: 1-16.
- EDMOND, D.B. (1966). - The influence of animal treading on pasture growth.
Proc. X. Int. Grassl. Cong., Finland, pp. 453-458.

- ELLENBERG, H. VON (1952). - "Landwirtschaftliche Pflanzensoziologie, II". Eugen Ulmer, Stuttgart, West Germany. p. 132.
- EVANS, P.S. (1967). - Leaf strength studies of pasture grasses. *J. Agric. Sci.* 69: 175-181.
- EWER, T.K. and SINCLAIR, D.P. (1952). - Lamb fattening trials on the Canterbury Plains. *N.Z. J. Sci. Technol.* 33A(6): 1-30.
- FELS, H.E., MOIR, R.J. and ROSSITER, R.C. (1959). - Herbage intake of grazing sheep in south-western Australia. *Aust. J. Agric. Res.* 10: 237-247.
- FITZGERALD, R.D. (1976). - Effect of stocking rate, lambing time and pasture management on wool and lamb production on annual subterranean clover pasture. *Aust. J. Agric. Res.* 27: 261-275.
- FRAME, J. (1971). - Fundamentals of grassland management, Part 10: The grazing animal. *Scot. Agric.* 50: 1-17.
- FRANKLIN, M.C. and POWNING, R.F. (1942). - Studies on the chemical composition of pods and seeds of certain species of *Medicago* with a note on the apparent digestibility of cluster clover (*Trifolium glomeratum* L.) seed. *J. Coun. Sci. Industr. Res. Aust.* 15: 191-200.
- GALLAGHER, J.R., WATKIN, B.R. and GRIMES, R.C. (1966). - An evaluation of pasture quality with young grazing sheep. 1. Live-weight growth and clean wool production. *J. Agric. Sci.* 66: 107-111.
- GRADWELL, M.W. (1966). - Soil moisture deficiencies in puddled pastures. *N.Z. J. Agric. Res.* 9: 127-136.
- GREENWOOD, E.A.N., DAVIES, H.L. and WATSON, E.R. (1967). - Growth of an annual pasture on virgin land in south-western Australia including effects of stocking rate and nitrogen fertilizer. *Aust. J. Agric. Res.* 18: 447-459.
- GUILBERT, H.R. and MEAD, S.W. (1931). - The digestibility of burr clover as affected by exposure to sunlight and rain. *Hilgardia* 6: 1-11.
- HAMILTON, B.A., HUTCHINSON, K.J. and SWAIN, F.G. (1970). - The growth of Merino lambs grazing four temperate grasses. *Proc. Aust. Soc. Anim. Prod.* 8: 455-459.
- HART, G.H., GUILBERT, H.R. and GOSS, H. (1932). - Seasonal changes in the chemical composition of range forage and their relation to nutrition of animals. California, Univ. Agric. Expt. Sta. Bull. No. 543, pp. 1-62.
- HEADY, H.F. (1961). - Continuous vs specialized grazing systems: a review and application to the California annual type. *J. Range Management* 14: 182-193.

- HEINEMANN, W.W. and VAN KEUREN, R.W. (1958). - A comparison of grass legume mixtures, legumes and grass under irrigation, as pastures for sheep.
Agron. J. 50: 189-192.
- HIGHT, G.K. and SINCLAIR, D.P. (1965). - The autumn growth of hoggets on different pasture types.
N.Z. J. Agric. Res. 8: 1079-1086.
- HILDER, E.J. (1964). - The distribution of plant nutrients by sheep at pasture.
Proc. Aust. Soc. Anim. Prod. 5: 241-248.
- HILDER, E.J. (1966). - Distribution of excreta by sheep at pasture.
Proc. X. Int. Grassl. Cong., Helsinki, Finland, Vol. 1, pp. 977-981.
- HODGE, R.W. and DOYLE, J.J. (1967). - Diet selected by lambs and yearling sheep grazing on annual and perennial pastures in southern Victoria.
Aust. J. Exp. Agric. Anim. Husb. 7: 141-143.
- HUTCHINGS, R.J., NEALE-SMITH, C.A. and SETON, R. (1963). - The behaviour under grazing of some introduced perennial grasses.
Fld. Stn. Rec. Div. Pl. Ind. CSIRO. (Aust.) 2(1) pp. 29-38.
- HUTCHINSON, J.C.D. and MANIKA WODZICKA-TOMASZEWSKA (1961). - Climate physiology of sheep.
Anim. Breed. Abstr. 29: 1-14.
- HUTCHINSON, K.J. (1970). - The persistence of perennial species under intensive grazing in a cool temperate environment.
Proc. XI. Int. Grassl. Cong., Qld. Aust. pp. 611-614.
- HUTCHINSON, K.J. and PORTER, R.B. (1958). - Growth and wool production of Merino hoggets related to grazing intake in a South Australian environment.
Proc. Aust. Soc. Anim. Prod. 2: 33-41.
- JOHNS, A.T. (1966). - Pasture content has dramatic effect on animal nutrition.
N.Z. J. Agric. 112: 44-47.
- JONES, D.B., CHRISTIAN, K.R. and SNAYDON, R.W. (1971). - Chemical composition and *in vitro* digestibility of some weed species during summer.
Aust. J. Exp. Agric. Anim. Husb. 11: 403-406.
- JONES, D.I.H. (1972). - The chemistry of grass for animal production.
Outlook on Agric. 7: 32-38.
- JONES, M.B. and EVANS, R.A. (1960). - Botanical composition changes in annual grassland as affected by fertilization and grazing.
Agron. J. 52: 459-461.

- JOYCE, J.P. and NEWTH, R.P. (1967). - Use of comparative slaughter technique to estimate the nutritive value of pasture for hoggets.
Proc. N.Z. Soc. Anim. Prod. 27: 166-180.
- KANWAR, J.S. (1956). - Phosphate retention in some Australian soils.
Soil Sci. 82: 43-50.
- KURTZ, L.T. and QUIRK, J.P. (1965). - Phosphate adsorption and phosphate fractions in field soils of varying history of phosphate fertilization.
Aust. J. Agric. Res. 16: 403-412.
- LAMBOURNE, L.J. (1969). - Evaluation of herbage plants by measurement of animal production.
Proc. Aust. Grassl. Cong. Perth, W. Aust. pp. 33-46.
- LANGLANDS, J.P. and BENNETT, I.L. (1973). - Stocking intensity and pastoral production. III. Wool production, fleece characteristics, and the utilization of nutrients for maintenance and wool growth by Merino sheep grazed at different stocking rates.
J. Agric. Sci. Camb. 81: 211-218.
- LIPSON, N. and BACON-HALL, R.E. (1974). - Influence of stocking rate on the manufacturing performance and textile properties of wool.
J. Aust. Inst. Agric. Sci. 40: 71-73.
- McARTHUR, I.D. (1970). - Optimum sheep stocking rate.
J. Aust. Inst. Agric. Sci. 36: 9-14.
- McARTHUR, I.D. and DILLON, J.L. (1971). - Risk, utility and stocking rate.
Aust. J. Agric. Econ. 15: 20-35.
- McINTYRE, G.A. (1946). - The measurement of pasture yield under grazing. Appendix in "Grazing management: continuous and rotational grazing by Merino sheep".
C.S.I.R. Aust. Bull. No. 201, pp. 70-83.
- McIVOR, J.G. and SMITH, D.F. (1973a). - A comparison of the effects of autumn deferment and continuous grazing on the growth of an annual pasture.
Aust. J. Exp. Agric. Anim. Husb. 13: 669-675.
- McIVOR, J.G. and SMITH, D.F. (1973b). - The effect of management during spring on the growth of a mixed annual pasture containing capeweed (*Arctotheca calendula*).
Aust. J. Exp. Agric. Anim. Husb. 13: 398-403.
- McIVOR, J.G. and SMITH, D.F. (1973c). - Plant factors influencing the nutritive value of some temperate annual pasture species.
Aust. J. Exp. Agric. Anim. Husb. 13: 404-410.

- McLEAN, J.W., THOMSON, G.C., IVERSEN, C.E., JAGUSH, K.T. and LAWSON, B.M. (1962). - Sheep production and health on pure species pastures.
Proc. N.Z. Grassld. Assoc. 24: 57-70.
- McMANUS, W.R., ARNOLD, G.W. and PAYNTER, J.R. (1964). - Studies in the wool production of grazing sheep. 2. Variation in wool characteristics with season and stocking rate.
Aust. J. Exp. Agric. Anim. Husb. 4: 404-411.
- McWILLIAM, J.R. (1962). - Interspecific hybridization in phalaris; hybrids between *Phalaris tuberosa* and the hexaploid race of *Phalaris arundinacea*.
Aust. J. Agric. Res. 13: 585-598.
- McWILLIAM, J.R., MACKAY, J.H.E., ROGERS, V.E. and SCHROEDER, H.E. (1965). - Characteristics and performance of interspecific hybrids between *Phalaris tuberosa* and *Phalaris arundinacea*. CSIRO. Aust. Div. Pl. Ind. Fld. Stn. Rec. 4, pp. 34-58.
- McWILLIAM, J.R. and NEAL-SMITH, C.A. (1962). - Tetraploid and hexaploid chromosome races of *Phalaris arundinacea* L.
Aust. J. Agric. Res. 13: 1-9.
- MANN, P.P., GORDDARD, B.J., GLENCROSS, R.N. and FITZPATRICK, E.N. (1966). - Stocking rate and rate of superphosphate in a higher rainfall area.
J. Dep. Agric. W. Aust. 7: 482-488.
- MINSON, D.J., RAYMOND, W.F. and HARRIS, C.E. (1960). - Studies in the digestibility of herbage. VIII. The digestibility of S37 cocksfoot, S23 ryegrass and S24 ryegrass.
J. Brit. Grassl. Soc. 15: 174-180.
- MOORE, R.M. (1965). - Ecological effects of grazing on grasslands in south-eastern Australia.
Proc. IX. Int. Grassl. Cong., Sao Paulo, Brazil. Vol. 1, pp. 429-433.
- MORLEY, F.H.W., BENNETT, D. and MCKINNEY, G.T. (1969). - The effect of intensity of rotational grazing with breeding ewes on phalaris-subterranean clover pastures.
Aust. J. Exp. Agric. Anim. Husb. 9: 74-84.
- MORLEY, F.H.W. and SPEDDING, C.R.W. (1968). - Agricultural systems and grazing experiments.
Herb. Abstr. 38: 279-287.
- MORRIS, L.R. (1961). - Photoperiodicity of seasonal rhythm of wool growth in sheep.
Nature 190: 102-103.
- MOWAT, D.N., FULKERSON, R.S., TOSSELL, W.E. and WINCH, J.E. (1965). - The *in vitro* dry matter digestibility of several species and varieties and their plant parts with advancing stages of maturity.
Proc. IX. Int. Grassl. Cong., Sao Paulo, Brazil. Vol. 1, pp. 801-806.

- MYERS, L.F. and SQUIRES, V.R. (1968a). - Some effects of cutting for hay on production of irrigated annual pastures in the following year.
Aust. J. Exp. Agric. Anim. Husb. 8: 197-202.
- MYERS, L.F. and SQUIRES, V.R. (1968b). - Comparison under grazing of *Phalaris tuberosa* and an interspecific *Phalaris* hybrid in irrigated swards.
Aust. Grassl. Conf. Perth, W. Aust. 1(2b), pp. 26-27.
- MYERS, L.F. and SQUIRES, V.R. (1970). - Control of barley grass by grazing management in irrigated pastures.
Aust. J. Exp. Agric. Anim. Husb. 10: 151-155.
- NEAL-SMITH, C.A. (1942). - The comparative grazing values of top-dressed natural and sown pastures in the middle south-east of South Australia.
J. Dep. Agric. S. Aust. 45: 485-496.
- NEAL-SMITH, C.A. (1955). - Report on herbage plant exploration in the Mediterranean region.
F.A.O. Rep. No. 415 (Rome).
- NORTHCOTE, K.H. and TUCKER, B.M. (1948). - A soil survey of the Hundred of Seddon and part of the Hundred of MacGillivray, Kangaroo Island, South Australia.
C.S.I.R. Aust. Bull. No. 233.
- PARROTT, R.T. and DONALD, C.M. (1970). - Growth and ignitability of annual pastures in a Mediterranean environment.
I. Effect of length of season and of defoliation on the growth, water content and desiccation of annual pastures.
Aust. J. Exp. Agric. Anim. Husb. 10: 67-75.
- PEARCE, G.R., VERCOE, J.E. and FREER, M. (1962). - The establishment of faecal nitrogen-digestibility regressions for animals grazing irrigated pasture.
J. Agric. Sci. 59: 397-402.
- PRITCHARD, G.I., FOLKINS, L.P. and PIGDEN, W.J. (1963). - The *in vitro* digestibility of whole grasses and their parts at progressive stages of maturity.
Can. J. Plant Sci. 43: 79-87.
- PULLMAN, A.L. and ALLDEN, W.G. (1971). - Chemical curing of annual pastures in southern Australia for beef cattle and sheep.
Aust. J. Agric. Res. 22: 401-413.
- RADCLIFFE, J.C. and COCHRANE, M.J. (1970). - Digestibility and crude protein changes in ten maturing pasture species.
Proc. Aust. Soc. Anim. Prod. 8: 531-536.

- RADCLIFFE, J.C. and NEWBERY, T.R. (1968). - Digestibility, crude protein and yield changes in maturing grass-legume hay crops.
Proc. Aust. Soc. Anim. Prod. 7: 66-69.
- RAE, A.L., BROUGHAM, R.W. and BARTON, R.A. (1964). - A note on the liveweight gains of sheep grazing different ryegrass pastures.
N.Z. J. Agric. Res. 7: 491-495.
- RAE, A.L., BROUGHAM, R.W., GLENDAY, A.C. and BUTLER, G.W. (1963). - Pasture type in relation to liveweight gain, carcass composition, iodine nutrition and some rumen characteristics of sheep. 1. Liveweight growth of the sheep.
J. Agric. Sci. Camb. 61: 187-190.
- RATCLIFF, R.D. and HEADY, H.F. (1962). - Seasonal changes in herbage weight in an annual grass community.
J. Range Management 15: 146-149.
- RAYMOND, W.F. (1969). - The nutritive value of forage crops.
Adv. Agron. 21: 1-108.
- REED, K.F.M. (1970). - Variation in liveweight gain with grass species in grass-clover pastures.
Proc. XI. Int. Grassl. Cong. Qld. Aust. pp. 877-880.
- REED, K.F.M. (1972a). - Some effects of botanical composition of pasture on growth and wool production of weaner sheep.
Aust. J. Exp. Agric. Anim. Husb. 12: 355-360.
- REED, K.F.M. (1972b). - The performance of sheep grazing different pasture types. In "Plants for sheep in Australia" pp. 193-204. A Review of pasture, browse and fodder crop research 1948-70. Eds. Leigh and Noble. (Angus and Robertson, Aust.).
- REED, K.F.M. (1974). - The productivity of pastures sown with *Phalaris tuberosa* or *Lolium perenne*.
1. Pasture growth and composition.
Aust. J. Exp. Agric. Anim. Husb. 14: 640-648.
- ROE, R., SOUTHCOTT, W.H. and TURNER, H.N. (1959). - Grazing management of pastures in the New England region of New South Wales.
1. Pasture and sheep production with special reference to systems of grazing and internal parasites.
Aust. J. Agric. Res. 10: 530-554.
- ROSSITER, R.C. (1958). - Utilization of annual pastures in a Mediterranean environment with special reference to autumn deferment.
Aust. Agrostology Conf., Univ. New England. Vol. 1, pp. 56/1-15.
- ROSSITER, R.C. (1961). - The influence of defoliation on the components of seed yield in swards of subterranean clover (*Trifolium subterraneum* L.).
Aust. J. Agric. Res. 12: 821-833.

- ROSSITER, R.C. (1966). - Ecology of the Mediterranean annual-type pasture.
Adv. Agron. 18: 1-56.
- ROSSITER, R.C. and PACK, R.J. (1956). - The effect of protection from grazing on the botanical composition of an annual-type subterranean clover pasture.
J. Aust. Inst. Agric. Sci. 22: 71-73.
- SHARKEY, M.J., DAVIS, I.F. and KENNEY, P.A. (1962). - The effect of previous and current nutrition on wool production in southern Victoria.
Aust. J. Exp. Agric. Anim. Husb. 2: 160-169.
- SHARKEY, M.J., DAVIS, I.F. and KENNEY, P.A. (1964). - The effect of rate of stocking with sheep on the botanical composition of an annual pasture in southern Victoria.
Aust. J. Exp. Agric. Anim. Husb. 4: 34-38.
- SHORT, B.F. and CHAPMAN, R.E. (1965). - Techniques for investigating wool growth. In "Field Investigations with Sheep - A Manual of Techniques", Section 13. Ed. G.R. Moule (CSIRO. Melb.), pp. 1-23.
- SILSBURY, J.H. (1961). - A study of dormancy, survival, and other characteristics in *Lolium perenne* L. at Adelaide, South Australia.
Aust. J. Agric. Res. 12: 1-9.
- SINCLAIR, D.P., CLARK, E.A. and FILMER, D.B. (1956). - A comparison of hogget ill-thrift on perennial ryegrass and white clover.
Proc. N.Z. Soc. Anim. Prod. 16: 23-28.
- SIVALINGAM (1973). - Some effects of treading by sheep on pastures in a Mediterranean environment of South Australia.
M.Ag.Sc. Thesis, University of Adelaide, Adelaide.
- SMITH, D.F. (1966). - Barley grass in Australian pastures.
Ph.D. Thesis, University of Melbourne, Melbourne.
- SMITH, D.F. (1968a). - The growth of barley grass (*Hordeum leporinum*) in annual pasture: 2. Growth comparisons with Wimmera ryegrass (*Lolium rigidum*).
Aust. J. Exp. Agric. Anim. Husb. 8: 484-490.
- SMITH, D.F. (1968b). - The growth of barley grass (*Hordeum leporinum*) in annual pasture: 4. The effect of some management practices on barley grass content.
Aust. J. Exp. Agric. Anim. Husb. 8: 706-711.
- SMITH, R.C.G., BIDDISCOMBE, E.F. and STERN, W.R. (1973). - Effect of spelling newly sown pastures.
Aust. J. Exp. Agric. Anim. Husb. 13: 549-555.
- SPEDDING, C.R.W., LARGE, R.V. and KYDD, D.D. (1966). - The evaluation of herbage species by grazing animals.
Proc. X. Int. Grassl. Cong., Helsinki, Finland. pp. 479-483.

- STEWART, A.M., MOIR, R.J. and SCHINCKEL, P.G. (1961). - Seasonal fluctuations in wool growth in south-western Australia. *Aust. J. Exp. Agric. Anim. Husb.* 6: 85-91.
- SUCKLING, F.E.T. (1952). - Dissemination of white clover by sheep. *N.Z. J. Sci. Tech.* 33: 64-77.
- TALBOT, M.W. and BISWELL, H.H. (1942). - The forage crop and its management. California, Univ. Agric. Expt. Sta. Bull. 663: 13-49.
- TAYLOR, G.B. (1972). - Time distribution of seedling emergence from single seed crops of several annual pasture legumes. *Aust. J. Exp. Agric. Anim. Husb.* 12: 628-633.
- TAYLOR, R.J. (1961). - Renovation is the answer to run-down pastures. *J. Dep. Agric. S. Aust.* 64: 510-518.
- TIVER, N.S. and CROCKER, R.L. (1951). - The grasslands of south east South Australia in relation to climate, soils and developmental history. *J. Brit. Grassl. Soc.* 6: 29-80.
- VAN DYNE, G.M. and HEADY, H.F. (1965). - Interrelations of botanical and chemical dietary components of animals grazing dry annual range. *J. Anim. Sci.* 24: 305-312.
- VERCOE, J.E. and PEARCE, G.R. (1960). - Digestibility of *Medicago tribuloides* (barrel medic) pods. *J. Aust. Inst. Agric. Sci.* 26: 67-70.
- WALKER, J. (1956). - The reaction of subterranean clover varieties to scorch caused by *Kabatiella caulivora* (Kirchn.) Karak. *J. Aust. Inst. Agric. Sci.* 22: 288-291.
- WEIR, W.C., MEYER, J.H. and LOFGREEN, G.P. (1959). - Symposium on forage evaluation: VI. The use of oesophageal fistula, lignin and chromogen techniques for studying selective grazing and digestibility of range and pasture by sheep and cattle. *Agron. J.* 51: 235-237.
- WHITE, D.H. and McCONCHIE, B.J. (1976). - Effect of stocking rate on fleece measurements and their relationships in Merino sheep. *Aust. J. Agric. Res.* 27: 163-174.
- WILSON, R.K. and McCARRICK, R.B. (1966). - Apparent dry-matter digestibility, voluntary food-intake and yields of dry matter of mixed swards, conserved as artificially dried grass and tetrapod hay, at progressive stages of maturity. Proc. X. Int. Grassl. Cong., Helsinki, Finland. pp. 371-379.